

BRNO UNIVERSITY OF TECHNOLOGY
VYSOKÉ UČENÍ TECHNICKÉ V BRNĚ

FACULTY OF ELECTRICAL ENGINEERING AND COMMUNICATION
DEPARTMENT OF MATHEMATICS

FAKULTA ELEKTROTECHNIKY A KOMUNIKAČNÍCH TECHNOLOGIÍ
ÚSTAV MATEMATIKY

WEAKLY DELAYED LINEAR PLANAR SYSTEMS OF
DISCRETE EQUATIONS

DOCTORAL THESIS
DIZERTAČNÍ PRÁCE

AUTHOR
AUTOR PRÁCE

Mgr. HANA HALFAROVÁ



BRNO UNIVERSITY OF TECHNOLOGY
VYSOKÉ UČENÍ TECHNICKÉ V BRNĚ



FACULTY OF ELECTRICAL ENGINEERING AND
COMMUNICATION
DEPARTMENT OF MATHEMATICS

FAKULTA ELEKTROTECHNIKY A KOMUNIKAČNÍCH
TECHNOLOGIÍ
ÚSTAV MATEMATIKY

WEAKLY DELAYED LINEAR PLANAR SYSTEMS OF
DISCRETE EQUATIONS
SLABĚ ZPOŽDĚNÉ LINEÁRNÍ ROVINNÉ SYSTÉMY DISKRÉTNÍCH ROVNIC

DOCTORAL THESIS
DIZERTAČNÍ PRÁCE

AUTHOR
AUTOR PRÁCE

Mgr. HANA HALFAROVÁ

SUPERVISOR
VEDOUCÍ PRÁCE

prof. RNDr. JOSEF DIBLÍK, DrSc.

BRNO 2014

ABSTRACT

The present thesis deals with planar weakly delayed linear discrete systems

$$x(k+1) = Ax(k) + \sum_{l=1}^n B^l x_l(k - m_l)$$

where $k \in \mathbb{Z}_0^\infty := \{0, 1, \dots, \infty\}$, m_1, m_2, \dots, m_n are constant integer delays, $0 < m_1 < m_2 < \dots < m_n$, A, B^1, \dots, B^n are constant 2×2 matrices and $x: \mathbb{Z}_{-m_n}^\infty \rightarrow \mathbb{R}^2$. The characteristic equations of weakly delayed systems are identical with those of the same systems but without delayed terms. In this case, after several steps, the space of solutions with a given starting dimension $2(m_n + 1)$ is pasted into a space with a dimension less than the starting one. In a sense, this situation is analogous to one known in the theory of linear differential systems with constant coefficients and special delays when the initially infinite dimensional space of solutions on the initial interval turns (after several steps) into a finite dimensional set of solutions. For every possible case, explicit general solutions are constructed and, finally, results on the dimensionality of the space of solutions are obtained. The stability of solutions is investigated as well.

KEYWORDS

discrete equation, linear systems of difference equations, weakly delayed system, planar system, dimension of the space of solutions, conditional stability

ABSTRAKT

Dizertační práce se zabývá slabě zpožděnými lineárními rovinnými systémy s konstantními koeficienty tvaru

$$x(k+1) = Ax(k) + \sum_{l=1}^n B^l x_l(k - m_l)$$

kde $k \in \mathbb{Z}_0^\infty := \{0, 1, \dots, \infty\}$, m_1, m_2, \dots, m_n jsou konstantní celá čísla, $0 < m_1 < m_2 < \dots < m_n$, A, B^1, \dots, B^n jsou konstantní 2×2 matice a $x: \mathbb{Z}_{-m_n}^\infty \rightarrow \mathbb{R}^2$ je hledané řešení. Charakteristická rovnice těchto systémů je identická s charakteristickou rovnicí systému, který neobsahuje zpožděné členy. V takovém případě se počáteční dimenze prostoru řešení $2(m_n + 1)$ mění po několika krocích na menší. V jistém smyslu je tato situace analogická podobnému jevu v teorii lineárních diferenciálních systémů s konstantními koeficienty a speciálním zpožděním, kdy původně nekonečně rozměrný prostor řešení (na počátečním intervalu) přejde po několika krocích do konečného prostoru řešení. V práci je pro každý možný případ kombinace kořenů charakteristické rovnice konstruováno obecné řešení daného systému a jsou formulovány výsledky o dimenzi prostoru řešení. Také je zkoumána stabilita řešení.

KLÍČOVÁ SLOVA

diskrétní rovnice, lineární systémy diferenčních rovnic, slabě zpožděný systém, rovinný systém, dimenze prostoru řešení, podmíněná stabilita

HALFAROVÁ, HANA *Weakly delayed linear planar systems of discrete equations*: doctoral thesis. Brno: Brno University of Technology, Faculty of Electrical Engineering and Communication, Department of Mathematics, 2014. 86 p. Supervised by prof. RNDr. Josef Diblík, DrSc.

DECLARATION

I declare that I have written my doctoral thesis on the theme of “Weakly delayed linear planar systems of discrete equations” independently, under the guidance of the doctoral thesis supervisor and using the technical literature and other sources of information which are all quoted in the thesis and detailed in the list of literature at the end of the thesis.

As the author of the doctoral thesis I furthermore declare that, as regards the creation of this doctoral thesis, I have not infringed any copyright. In particular, I have not unlawfully encroached on anyone’s personal and/or ownership rights and I am fully aware of the consequences in the case of breaking Regulation § 11 and the following of the Copyright Act No 121/2000 Sb., and of the rights related to intellectual property right and changes in some Acts (Intellectual Property Act) and formulated in later regulations, inclusive of the possible consequences resulting from the provisions of Criminal Act No 40/2009 Sb., Section 2, Head VI, Part 4.

Brno

.....

(author’s signature)

ACKNOWLEDGEMENT

I would like to sincerely thank my supervisor prof. RNDr. Josef Diblík, DrSc. for his helpful advice, lots of worthy suggestions and great patience.

Brno

.....

(author's signature)

CONTENTS

1	Introduction	8
1.1	Current State	9
1.2	Aims of the thesis	9
1.3	Preliminary notions and properties	10
1.4	Weakly delayed systems	10
1.5	Necessary and sufficient conditions determining weakly delayed systems	12
1.6	Problem under consideration	17
1.7	Auxiliary formula	17
2	General solution	18
2.1	Jordan forms of the matrix \mathbf{A} and corresponding solutions of the problem (1.1), (1.4)	18
2.1.1	The case (2.2) of two real distinct roots	19
2.1.2	The case (2.3) of two complex conjugate roots	28
2.1.3	The case (2.4) of double real root	28
2.1.4	The case (2.5) of a double real root	56
3	Dimension of the set of solutions	65
4	Discussion of stability	69
5	Conclusions	82
	References	83

1 INTRODUCTION

As most of the time measurements involving variables are discrete, the observed evolution phenomena can be expressed naturally in terms of difference equations and, thus, such equations are important mathematical models in their own right. Difference equations have an important role in the study of discretization methods for differential equations, too. The theory of difference equations is much richer than the corresponding theory of differential equations. For example, a simple difference equation obtained from a first-order differential equation may involve phenomena that can only occur for higher-order differential equations. Thus, the theory of difference equations is interesting by itself and, therefore, likely to take on greater importance in the near future.

The application of the theory of difference equations is rapidly increasing in various fields such as numerical analysis, control theory, finite mathematics, and computer science.

The fundamentals of the theory of difference equations are well described for example in books by S. Elaydi [19], by I. Györi, G. Ladas [21], by V. L. Kocić, G. Ladas [27] and by R. P. Agarwal, M. Bohner, S. R. Grace, D. O'Regan [1].

1.1 Current State

The difference equations have recently been an object of intensive research. Monographs summarizing some outcomes were mentioned above.

Every month, numerous new papers are published on the qualitative theory of difference equations. Some interesting results have been published on the representations of solutions of linear discrete systems with delay, e.g., [13, 14, 25], on the existence of positive solutions of discrete equations, e.g., [2, 3, 4, 5, 29], on the oscillation of solutions of discrete equations, e.g., [23, 30, 31], on the stability of solutions of discrete systems, e.g., [6, 16, 22, 26, 28] and on the asymptotic properties of solutions of the discrete equations and systems such as [7, 8, 11, 12, 17, 18, 20, 39].

1.2 Aims of the thesis

The thesis is concened with planar systems of weakly delayed equations

$$x(k+1) = Ax(k) + \sum_{l=1}^n B^l x_l(k - m_l) \quad (1.1)$$

where m_1, m_2, \dots, m_n are constant integer delays, $0 < m_1 < m_2 < \dots < m_n$, $k \in \mathbb{Z}_0^\infty$, A, B^1, \dots, B^n are constant 2×2 matrices, $A = (a_{ij})$, $B^l = (b_{ij}^l)$, $i, j = 1, 2$, $l = 1, 2, \dots, n$ and $x: \mathbb{Z}_{-m_n}^\infty \rightarrow \mathbb{R}^2$. In the thesis, we construct general solutions of such systems. Further, we show that, after several steps, the dimension of the space of all solutions is reduced to a less-dimensional space. Moreover, we discuss the stability of the system.

Methodically, we follow the paper [15] where a planar weakly delayed linear discrete system

$$x(k+1) = Ax(k) + Bx(k-m), \quad (1.2)$$

is considered with $m \geq 0$ being a fixed integer, $k \in \mathbb{Z}_0^\infty$, $A = (a_{ij})$, $B = (b_{ij})$ constant 2×2 matrices, and $x: \mathbb{Z}_{-m}^\infty \rightarrow \mathbb{R}^2$. A general solution of (1.2) is constructed and results on the dimensionality of the space of solutions are derived.

Some of the results obtained are published in [9, 10]. In [9], a system (1.1) with $n = 2$ is considered. The results published in [10] concern system (1.1) and generalize the results published in [15].

1.3 Preliminary notions and properties

We use the following notation: for integers s, q , $s \leq q$, we define $\mathbb{Z}_s^q := \{s, s + 1, \dots, q\}$ where $s = -\infty$ or $q = \infty$ are admitted, too. Throughout this dissertation, using notation \mathbb{Z}_s^q , we always assume $s \leq q$. In the thesis, we deal with the discrete planar systems (1.1)

$$x(k+1) = Ax(k) + \sum_{l=1}^n B^l x_l(k - m_l)$$

where m_1, m_2, \dots, m_n are constant integer delays, $0 < m_1 < m_2 < \dots < m_n$, $k \in \mathbb{Z}_0^\infty$, A, B^1, \dots, B^n are constant 2×2 matrices, $A = (a_{ij})$, $B^l = (b_{ij}^l)$, $i, j = 1, 2$, $l = 1, 2, \dots, n$ and $x: \mathbb{Z}_{-m_n}^\infty \rightarrow \mathbb{R}^2$. Throughout the dissertation, we assume that

$$B^l \neq \Theta \tag{1.3}$$

where $l = 1, 2, \dots, n$ and Θ is 2×2 zero matrix. Together with equation (1.1), we consider an initial (Cauchy) problem

$$x(k) = \varphi(k) \tag{1.4}$$

where $k = -m_n, -m_n + 1, \dots, 0$ with $\varphi: \mathbb{Z}_{-m_n}^0 \rightarrow \mathbb{R}^2$. The *existence* and *uniqueness* of the solution of the initial problem (1.1), (1.4) on $\mathbb{Z}_{-m_n}^\infty$ is obvious. We recall that the *solution* $x: \mathbb{Z}_{-m_n}^\infty \rightarrow \mathbb{R}^2$ of (1.1), (1.4) is defined as an *infinite sequence*

$$\{x(-m_n) = \varphi(-m_n), x(-m_n + 1) = \varphi(-m_n + 1), \dots, \\ x(0) = \varphi(0), x(1), x(2), \dots, x(k), \dots\}$$

such that, for any $k \in \mathbb{Z}_0^\infty$, equality (1.1) holds.

The space of all initial data (1.4) with $\varphi: \mathbb{Z}_{-m_n}^0 \rightarrow \mathbb{R}^2$ is obviously $2(m_n + 1)$ -dimensional. Below, we describe the fact that, among the systems (1.1), there are such systems that their space of solutions, being initially $2(m_n + 1)$ -dimensional, on a reduced interval turns into a space having a dimension less than $2(m_n + 1)$. The problem under consideration (pasting property of solutions) is exactly formulated in Part 1.4.

1.4 Weakly delayed systems

We consider the system (1.1) and look for a solution having the form $x(k) = \xi \lambda^k$ where $k \in \mathbb{Z}_{-m_n}^\infty$, $\lambda = \text{const}$ with $\lambda \neq 0$ and $\xi = (\xi_1, \xi_2)^T$ is a nonzero constant vector. The usual procedure leads to a characteristic equation

$$D := \det \left(A + \sum_{l=1}^n \lambda^{-m_l} B^l - \lambda I \right) = 0 \tag{1.5}$$

where I is the unit 2×2 matrix. Together with (1.1), we consider a system with the terms containing delays omitted

$$x(k+1) = Ax(k) \quad (1.6)$$

and its characteristic equation

$$\det(A - \lambda I) = 0. \quad (1.7)$$

Definition 1.4.1. *The system (1.1) is called a weakly delayed system if the characteristic equations (1.5), (1.7) corresponding to systems (1.1) and (1.6) are equal, i.e. if, for every $\lambda \in \mathbb{C} \setminus \{0\}$,*

$$D = \det \left(A + \sum_{l=1}^n \lambda^{-m_l} B^l - \lambda I \right) = \det(A - \lambda I). \quad (1.8)$$

We consider a linear transformation

$$x(k) = \mathcal{S}y(k) \quad (1.9)$$

with a nonsingular 2×2 matrix \mathcal{S} . Then, the discrete system for y is

$$y(k+1) = A_S y(k) + \sum_{l=1}^n B_S^l y(k - m_l) \quad (1.10)$$

with $A_S = \mathcal{S}^{-1}A\mathcal{S}$, $B_S^l = \mathcal{S}^{-1}B^l\mathcal{S}$ where $l = 1, 2, \dots, n$. We show that a system's property of being one weakly delayed is preserved by every nonsingular linear transformation.

Lemma 1.4.2. *If the system (1.1) is a weakly delayed system, then its arbitrary linear nonsingular transformation (1.9) again leads to a weakly delayed system (1.10).*

Proof. It is easy to show that

$$\det \left(A_S + \sum_{l=1}^n \lambda^{-m_l} B_S^l - \lambda I \right) = \det(A_S - \lambda I)$$

holds since

$$\begin{aligned} & \det \left(A_S + \sum_{l=1}^n \lambda^{-m_l} B_S^l - \lambda I \right) \\ &= \det \left(A_S + \lambda^{-m_1} B_S^1 + \lambda^{-m_2} B_S^2 + \dots + \lambda^{-m_n} B_S^n - \lambda I \right) \\ &= \det \left(\mathcal{S}^{-1}A\mathcal{S} + \lambda^{-m_1} \mathcal{S}^{-1}B^1\mathcal{S} + \lambda^{-m_2} \mathcal{S}^{-1}B^2\mathcal{S} + \dots \right. \\ & \quad \left. + \lambda^{-m_n} \mathcal{S}^{-1}B^n\mathcal{S} - \lambda \mathcal{S}^{-1}I\mathcal{S} \right) \\ &= \det \left[\mathcal{S}^{-1} \left(A + \lambda^{-m_1} B^1 + \lambda^{-m_2} B^2 + \dots + \lambda^{-m_n} B^n - \lambda I \right) \mathcal{S} \right] \end{aligned}$$

$$\begin{aligned}
 &= \det \mathcal{S}^{-1} \det \left(A + \lambda^{-m_1} B^1 + \lambda^{-m_2} B^2 + \cdots + \lambda^{-m_n} B^n - \lambda I \right) \det \mathcal{S} \\
 &= \det \left(A + \lambda^{-m_1} B^1 + \lambda^{-m_2} B^2 + \cdots + \lambda^{-m_n} B^n - \lambda I \right) \\
 &= \det \left(A + \sum_{l=1}^n \lambda^{-m_l} B^l - \lambda I \right),
 \end{aligned}$$

$$\begin{aligned}
 \det (A_{\mathcal{S}} - \lambda I) &= \det \left(\mathcal{S}^{-1} A \mathcal{S} - \lambda \mathcal{S}^{-1} I \mathcal{S} \right) \\
 &= \det \left[\mathcal{S}^{-1} (A - \lambda I) \mathcal{S} \right] \\
 &= \det \mathcal{S}^{-1} \det (A - \lambda I) \det \mathcal{S} \\
 &= \det (A - \lambda I)
 \end{aligned}$$

and the equality

$$\det \left(A + \sum_{l=1}^n \lambda^{-m_l} B^l - \lambda I \right) = \det (A - \lambda I),$$

i.e. the equality (1.8) is assumed. □

1.5 Necessary and sufficient conditions determining weakly delayed systems

In the below theorem, we give conditions, in terms of determinants, indicating whether a system is weakly delayed.

Theorem 1.5.1. *System (1.1) is a weakly delayed system if and only if the following $3n + n(n-1)/2$ conditions hold simultaneously:*

$$b_{11}^l + b_{22}^l = 0, \tag{1.11}$$

$$\begin{vmatrix} b_{11}^l & b_{12}^l \\ b_{21}^l & b_{22}^l \end{vmatrix} = 0, \tag{1.12}$$

$$\begin{vmatrix} a_{11} & a_{12} \\ b_{21}^l & b_{22}^l \end{vmatrix} + \begin{vmatrix} b_{11}^l & b_{12}^l \\ a_{21} & a_{22} \end{vmatrix} = 0, \tag{1.13}$$

$$\begin{vmatrix} b_{11}^l & b_{12}^l \\ b_{21}^v & b_{22}^v \end{vmatrix} + \begin{vmatrix} b_{11}^v & b_{12}^v \\ b_{21}^l & b_{22}^l \end{vmatrix} = 0 \tag{1.14}$$

where $l, v = 1, 2, \dots, n$ and $v > l$.

Proof. We start with computing determinant D defined by (1.5). We get

$$D = \begin{vmatrix} D_1 & D_2 \\ D_3 & D_4 \end{vmatrix},$$

where

$$D_1 = a_{11} + b_{11}^1 \lambda^{-m_1} + b_{11}^2 \lambda^{-m_2} + \dots + b_{11}^n \lambda^{-m_n} - \lambda,$$

$$D_2 = a_{12} + b_{12}^1 \lambda^{-m_1} + b_{12}^2 \lambda^{-m_2} + \dots + b_{12}^n \lambda^{-m_n},$$

$$D_3 = a_{21} + b_{21}^1 \lambda^{-m_1} + b_{21}^2 \lambda^{-m_2} + \dots + b_{21}^n \lambda^{-m_n},$$

$$D_4 = a_{22} + b_{22}^1 \lambda^{-m_1} + b_{22}^2 \lambda^{-m_2} + \dots + b_{22}^n \lambda^{-m_n} - \lambda.$$

Expanding the determinant on the right-hand side along summands of the first column, we get:

$$\begin{aligned} D &= \begin{vmatrix} a_{11} & a_{12} + b_{12}^1 \lambda^{-m_1} + b_{12}^2 \lambda^{-m_2} + \dots + b_{12}^n \lambda^{-m_n} \\ a_{21} & a_{22} + b_{22}^1 \lambda^{-m_1} + b_{22}^2 \lambda^{-m_2} + \dots + b_{22}^n \lambda^{-m_n} - \lambda \end{vmatrix} \\ &+ \lambda^{-m_1} \begin{vmatrix} b_{11}^1 & a_{12} + b_{12}^1 \lambda^{-m_1} + b_{12}^2 \lambda^{-m_2} + \dots + b_{12}^n \lambda^{-m_n} \\ b_{21}^1 & a_{22} + b_{22}^1 \lambda^{-m_1} + b_{22}^2 \lambda^{-m_2} + \dots + b_{22}^n \lambda^{-m_n} - \lambda \end{vmatrix} \\ &+ \lambda^{-m_2} \begin{vmatrix} b_{11}^2 & a_{12} + b_{12}^1 \lambda^{-m_1} + b_{12}^2 \lambda^{-m_2} + \dots + b_{12}^n \lambda^{-m_n} \\ b_{21}^2 & a_{22} + b_{22}^1 \lambda^{-m_1} + b_{22}^2 \lambda^{-m_2} + \dots + b_{22}^n \lambda^{-m_n} - \lambda \end{vmatrix} \\ &+ \dots \\ &+ \lambda^{-m_n} \begin{vmatrix} b_{11}^n & a_{12} + b_{12}^1 \lambda^{-m_1} + b_{12}^2 \lambda^{-m_2} + \dots + b_{12}^n \lambda^{-m_n} \\ b_{21}^n & a_{22} + b_{22}^1 \lambda^{-m_1} + b_{22}^2 \lambda^{-m_2} + \dots + b_{22}^n \lambda^{-m_n} - \lambda \end{vmatrix} \\ &+ \lambda \begin{vmatrix} -1 & a_{12} + b_{12}^1 \lambda^{-m_1} + b_{12}^2 \lambda^{-m_2} + \dots + b_{12}^n \lambda^{-m_n} \\ 0 & a_{22} + b_{22}^1 \lambda^{-m_1} + b_{22}^2 \lambda^{-m_2} + \dots + b_{22}^n \lambda^{-m_n} - \lambda \end{vmatrix}. \end{aligned}$$

Expanding each of the above determinants along summands of the second column, we have

$$\begin{aligned} D &= \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix} + \lambda^{-m_1} \begin{vmatrix} a_{11} & b_{12}^1 \\ a_{21} & b_{22}^1 \end{vmatrix} + \lambda^{-m_2} \begin{vmatrix} a_{11} & b_{12}^2 \\ a_{21} & b_{22}^2 \end{vmatrix} \\ &+ \dots + \lambda^{-m_n} \begin{vmatrix} a_{11} & b_{12}^n \\ a_{21} & b_{22}^n \end{vmatrix} + \lambda \begin{vmatrix} a_{11} & 0 \\ a_{21} & -1 \end{vmatrix} \end{aligned}$$

$$\begin{aligned}
 & + \lambda^{-m_1} \left[\begin{vmatrix} b_{11}^1 & a_{12} \\ b_{21}^1 & a_{22} \end{vmatrix} + \lambda^{-m_1} \begin{vmatrix} b_{11}^1 & b_{12}^1 \\ b_{21}^1 & b_{22}^1 \end{vmatrix} + \lambda^{-m_2} \begin{vmatrix} b_{11}^1 & b_{12}^2 \\ b_{21}^1 & b_{22}^2 \end{vmatrix} \right. \\
 & + \dots + \lambda^{-m_n} \begin{vmatrix} b_{11}^1 & b_{12}^n \\ b_{21}^1 & b_{22}^n \end{vmatrix} + \lambda \begin{vmatrix} b_{11}^1 & 0 \\ b_{21}^1 & -1 \end{vmatrix} \left. \right] \\
 & + \lambda^{-m_2} \left[\begin{vmatrix} b_{11}^2 & a_{12} \\ b_{21}^2 & a_{22} \end{vmatrix} + \lambda^{-m_1} \begin{vmatrix} b_{11}^2 & b_{12}^1 \\ b_{21}^2 & b_{22}^1 \end{vmatrix} + \lambda^{-m_2} \begin{vmatrix} b_{11}^2 & b_{12}^2 \\ b_{21}^2 & b_{22}^2 \end{vmatrix} \right. \\
 & + \dots + \lambda^{-m_n} \begin{vmatrix} b_{11}^2 & b_{12}^n \\ b_{21}^2 & b_{22}^n \end{vmatrix} + \lambda \begin{vmatrix} b_{11}^2 & 0 \\ b_{21}^2 & -1 \end{vmatrix} \left. \right] \\
 & + \dots \\
 & + \lambda^{-m_n} \left[\begin{vmatrix} b_{11}^n & a_{12} \\ b_{21}^n & a_{22} \end{vmatrix} + \lambda^{-m_1} \begin{vmatrix} b_{11}^n & b_{12}^1 \\ b_{21}^n & b_{22}^1 \end{vmatrix} + \lambda^{-m_2} \begin{vmatrix} b_{11}^n & b_{12}^2 \\ b_{21}^n & b_{22}^2 \end{vmatrix} \right. \\
 & + \dots + \lambda^{-m_n} \begin{vmatrix} b_{11}^n & b_{12}^n \\ b_{21}^n & b_{22}^n \end{vmatrix} + \lambda \begin{vmatrix} b_{11}^n & 0 \\ b_{21}^n & -1 \end{vmatrix} \left. \right] \\
 & + \lambda \left[\begin{vmatrix} -1 & a_{12} \\ 0 & a_{22} \end{vmatrix} + \lambda^{-m_1} \begin{vmatrix} -1 & b_{12}^1 \\ 0 & b_{22}^1 \end{vmatrix} + \lambda^{-m_2} \begin{vmatrix} -1 & b_{12}^2 \\ 0 & b_{22}^2 \end{vmatrix} \right. \\
 & + \dots + \lambda^{-m_n} \begin{vmatrix} -1 & b_{12}^n \\ 0 & b_{22}^n \end{vmatrix} + \lambda \begin{vmatrix} -1 & 0 \\ 0 & -1 \end{vmatrix} \left. \right].
 \end{aligned}$$

After simplification, we get

$$\begin{aligned}
 D & = \begin{vmatrix} a_{11} - \lambda & a_{12} \\ a_{21} & a_{22} - \lambda \end{vmatrix} \\
 & - \lambda^{-m_1+1}(b_{11}^1 + b_{22}^1) - \lambda^{-m_2+1}(b_{11}^2 + b_{22}^2) + \dots - \lambda^{-m_n+1}(b_{11}^n + b_{22}^n) \\
 & + \lambda^{-m_1} \left[\begin{vmatrix} a_{11} & a_{12} \\ b_{21}^1 & b_{22}^1 \end{vmatrix} + \begin{vmatrix} b_{11}^1 & b_{12}^1 \\ a_{21} & a_{22} \end{vmatrix} \right] + \lambda^{-m_2} \left[\begin{vmatrix} a_{11} & a_{12} \\ b_{21}^2 & b_{22}^2 \end{vmatrix} + \begin{vmatrix} b_{11}^2 & b_{12}^2 \\ a_{21} & a_{22} \end{vmatrix} \right] \\
 & + \dots + \lambda^{-m_n} \left[\begin{vmatrix} a_{11} & a_{12} \\ b_{21}^n & b_{22}^n \end{vmatrix} + \begin{vmatrix} b_{11}^n & b_{12}^n \\ a_{21} & a_{22} \end{vmatrix} \right]
 \end{aligned}$$

$$\begin{aligned}
 & + \lambda^{-m_1-m_2} \left[\begin{vmatrix} b_{11}^1 & b_{12}^1 \\ b_{21}^2 & b_{22}^2 \end{vmatrix} + \begin{vmatrix} b_{11}^2 & b_{12}^2 \\ b_{21}^1 & b_{22}^1 \end{vmatrix} \right] + \cdots + \lambda^{-m_1-m_n} \left[\begin{vmatrix} b_{11}^1 & b_{12}^1 \\ b_{21}^n & b_{22}^n \end{vmatrix} + \begin{vmatrix} b_{11}^n & b_{12}^n \\ b_{21}^1 & b_{22}^1 \end{vmatrix} \right] \\
 & + \lambda^{-m_2-m_3} \left[\begin{vmatrix} b_{11}^2 & b_{12}^2 \\ b_{21}^3 & b_{22}^3 \end{vmatrix} + \begin{vmatrix} b_{11}^3 & b_{12}^3 \\ b_{21}^2 & b_{22}^2 \end{vmatrix} \right] + \cdots + \lambda^{-m_2-m_n} \left[\begin{vmatrix} b_{11}^2 & b_{12}^2 \\ b_{21}^n & b_{22}^n \end{vmatrix} + \begin{vmatrix} b_{11}^n & b_{12}^n \\ b_{21}^2 & b_{22}^2 \end{vmatrix} \right] \\
 & + \cdots + \lambda^{-m_{n-1}-m_n} \left[\begin{vmatrix} b_{11}^{n-1} & b_{12}^{n-1} \\ b_{21}^n & b_{22}^n \end{vmatrix} + \begin{vmatrix} b_{11}^n & b_{12}^n \\ b_{21}^{n-1} & b_{22}^{n-1} \end{vmatrix} \right] \\
 & + \lambda^{-2m_1} \begin{vmatrix} b_{11}^1 & b_{12}^1 \\ b_{21}^1 & b_{22}^1 \end{vmatrix} + \lambda^{-2m_2} \begin{vmatrix} b_{11}^2 & b_{12}^2 \\ b_{21}^2 & b_{22}^2 \end{vmatrix} + \cdots + \lambda^{-2m_n} \begin{vmatrix} b_{11}^n & b_{12}^n \\ b_{21}^n & b_{22}^n \end{vmatrix}.
 \end{aligned}$$

Now we see that, for (1.8) to hold, i.e.,

$$D = \det \left(A + \sum_{l=1}^n \lambda^{-m_l} B^l - \lambda I \right) = \det (A - \lambda I) = \begin{vmatrix} a_{11} - \lambda & a_{12} \\ a_{21} & a_{22} - \lambda \end{vmatrix}$$

conditions (1.11)–(1.14) are both necessary and sufficient. □

Lemma 1.5.2. *Conditions (1.11)–(1.14) are equivalent to*

$$\operatorname{tr} B^l = \det B^l = 0, \quad (1.15)$$

$$\det(A + B^l) = \det A, \quad (1.16)$$

$$\det(B^l + B^v) = 0, \quad (1.17)$$

where $l, v = 1, 2, \dots, n$ and $v > l$.

Proof. I. We show that assumptions (1.11)–(1.14) imply (1.15)–(1.17). It is obvious that condition (1.15) is equivalent to (1.11), (1.12). Now we consider

$$\det(A + B^l) = \begin{vmatrix} a_{11} + b_{11}^l & a_{12} + b_{12}^l \\ a_{21} + b_{21}^l & a_{22} + b_{22}^l \end{vmatrix}.$$

Expanding the determinant on the right-hand side along summands of the first column and then expanding each of the determinants along summands of the second column, we have

$$\det(A + B^l) = \begin{vmatrix} a_{11} & a_{12} + b_{12}^l \\ a_{21} & a_{22} + b_{22}^l \end{vmatrix} + \begin{vmatrix} b_{11}^l & a_{12} + b_{12}^l \\ b_{21}^l & a_{22} + b_{22}^l \end{vmatrix}$$

$$= \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix} + \begin{vmatrix} a_{11} & b'_{12} \\ a_{21} & b'_{22} \end{vmatrix} + \begin{vmatrix} b^l_{11} & a_{12} \\ b^l_{21} & a_{22} \end{vmatrix} + \begin{vmatrix} b^l_{11} & b^l_{12} \\ b^l_{21} & b^l_{22} \end{vmatrix} \quad (1.18)$$

$$= [\text{due to (1.12) and (1.13)}] = \det A.$$

Now we consider

$$\det(B^l + B^v) = \begin{vmatrix} b^l_{11} + b^v_{11} & b^l_{12} + b^v_{12} \\ b^l_{21} + b^v_{21} & b^l_{22} + b^v_{22} \end{vmatrix}.$$

Expanding the determinant on the right-hand side along summands of the first column and then expanding each of the determinants along summands of the second column, we have

$$\begin{aligned} \det(B^l + B^v) &= \begin{vmatrix} b^l_{11} & b^l_{12} + b^v_{12} \\ b^l_{21} & b^l_{22} + b^v_{22} \end{vmatrix} + \begin{vmatrix} b^v_{11} & b^l_{12} + b^v_{12} \\ b^v_{21} & b^l_{22} + b^v_{22} \end{vmatrix} \\ &= \begin{vmatrix} b^l_{11} & b^l_{12} \\ b^l_{21} & b^l_{22} \end{vmatrix} + \begin{vmatrix} b^l_{11} & b^v_{12} \\ b^l_{21} & b^v_{22} \end{vmatrix} + \begin{vmatrix} b^v_{11} & b^l_{12} \\ b^v_{21} & b^l_{22} \end{vmatrix} + \begin{vmatrix} b^v_{11} & b^v_{12} \\ b^v_{21} & b^v_{22} \end{vmatrix} \quad (1.19) \\ &= [\text{due to (1.12), (1.14)}] = 0. \end{aligned}$$

II. Now we prove that assumptions (1.15)–(1.17) imply (1.11) and (1.14). Due to equivalence (1.11)–(1.12) with (1.15), it remains to be shown that (1.15)–(1.17) imply (1.13) and (1.14).

If (1.16) holds, then, from computations (1.18), we see that

$$\begin{vmatrix} a_{11} & b^l_{12} \\ a_{21} & b^l_{22} \end{vmatrix} + \begin{vmatrix} b^l_{11} & a_{12} \\ b^l_{21} & a_{22} \end{vmatrix} + \begin{vmatrix} b^l_{11} & b^l_{12} \\ b^l_{21} & b^l_{22} \end{vmatrix} = 0,$$

and because of (1.15) we get (1.13).

Finally, we show that (1.15) and (1.17) implies (1.14). From (1.19), (using (1.15)), we get

$$\det(B^l + B^v) = \begin{vmatrix} b^l_{11} & b^v_{12} \\ b^l_{21} & b^v_{22} \end{vmatrix} + \begin{vmatrix} b^v_{11} & b^l_{12} \\ b^v_{21} & b^l_{22} \end{vmatrix} = 0,$$

i.e., (1.14) holds.

□

1.6 Problem under consideration

The aim of this thesis is to give explicit formulas for the solutions of weakly delayed systems. This is done in Chapter 2. Moreover, we show (in Chapter 3) that, after several steps, the dimension of the space of all solutions, being initially equal to the dimension $2(m_n + 1)$ of the space of initial data (1.4) generated by discrete functions φ , is reduced to a dimension less than the initial one on an interval of the form \mathbb{Z}_s^∞ with $s > 0$. In other words, we will show that the $2(m_n + 1)$ -dimensional space of all solutions of (1.1) is pasted to a less-dimensional space of solutions on \mathbb{Z}_s^∞ . This problem is solved directly by explicitly computing the corresponding solutions of the Cauchy problems with each of the cases arising being considered. The underlying idea for such investigation is simple. If (1.1) is a weakly delayed system, then the corresponding characteristic equation has only two eigenvalues instead of $2(m_n + 1)$ eigenvalues in the case of systems with non-weak delays. This explains why the dimension of the space of solutions becomes less than the initial one. The final results (Theorems 3.0.6 – 3.0.9 below) provide the dimension of the space of solutions. Our results (published in [9] and [10]) generalize the results in [15]. Paper [9] considers the system (1.1) with $n = 2$ and, in [10], a general case (1.1) is treated. From explicit formulas we deduce (in Chapter 4) stability and so-called conditional stability of the system (1.1).

1.7 Auxiliary formula

For the reader's convenience, we recall one explicit formula (see e.g. [19]) for the solutions of linear scalar discrete non-delayed equations used in this thesis. We consider initial - value problem for the first order linear discrete nonhomogeneous equation

$$w(k + 1) = aw(k) + g(k), \quad w(k_0) = w_0, \quad k \in \mathbb{Z}_{k_0}^\infty$$

with $a \in \mathbb{C}$ and $g: \mathbb{Z}_{k_0}^\infty \rightarrow \mathbb{C}$. Then, it is easy to verify that unique solution of this problem is

$$w(k) = a^{k-k_0} w_0 + \sum_{r=k_0}^{k-1} a^{k-1-r} g(r), \quad k \in \mathbb{Z}_{k_0+1}^\infty. \quad (1.20)$$

Throughout this thesis, we adopt the customary notation for the sum: $\sum_{i=\ell+t}^\ell \mathcal{F}(i) = 0$ where ℓ is an integer, t is a positive integer and “ \mathcal{F} ” denotes the function considered independently of whether it is defined for indicated arguments or not.

Note that the formula (1.20) is many times used in recent literature to analyze asymptotic properties of solutions of various classes of difference equations, including nonlinear equations. We refer, e.g., to [32]– [38] and to relevant references therein.

2 GENERAL SOLUTION

In this chapter we derive general solution of weakly delayed system (1.1). If (1.8) holds, then equations (1.5) and (1.7) have only two (and the same) roots simultaneously. In order to prove the properties of the family of solutions of (1.1) formulated in Introduction, we will discuss each combination of roots, i.e., the cases of two real and distinct roots, a pair of complex conjugate roots, and, finally, a double real root.

Although computations in Parts 1.4 and 1.5 were performed under assumption $\lambda \neq 0$, results of this part remain valid also if one or both roots of characteristic equation (1.7) are zero.

2.1 Jordan forms of the matrix A and corresponding solutions of the problem (1.1), (1.4)

It is known that, for every matrix A , there exists a nonsingular matrix S transforming it to the corresponding Jordan matrix form Λ . This means that

$$\Lambda = S^{-1}AS,$$

where Λ has the following four possible forms (denoted below as $\Lambda_1, \Lambda_2, \Lambda_3, \Lambda_4$), depending on the roots of the characteristic equation (1.7), i.e. on the roots of

$$\lambda^2 - (a_{11} + a_{22})\lambda + (a_{11}a_{22} - a_{12}a_{21}) = 0. \quad (2.1)$$

If (2.1) has two real distinct roots λ_1, λ_2 , then

$$\Lambda_1 = \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix}, \quad (2.2)$$

if the roots are complex conjugate, i.e. $\lambda_{1,2} = p \pm iq$ with $q \neq 0$, then

$$\Lambda_2 = \begin{pmatrix} p & q \\ -q & p \end{pmatrix} \quad (2.3)$$

and, finally, in the case of one double real root $\lambda_{1,2} = \lambda$, we have either

$$\Lambda_3 = \begin{pmatrix} \lambda & 0 \\ 0 & \lambda \end{pmatrix} \quad (2.4)$$

or

$$\Lambda_4 = \begin{pmatrix} \lambda & 1 \\ 0 & \lambda \end{pmatrix}. \quad (2.5)$$

The transformation $y(k) = S^{-1}x(k)$ transforms (1.1) into a system

$$y(k+1) = \Lambda y(k) + \sum_{l=1}^n B^{*l} y(k - m_l), \quad k \in \mathbb{Z}_0^\infty \quad (2.6)$$

with $B^{*l} = S^{-1}B^lS$, $B^{*l} = (b_{ij}^{*l})$, $l = 1, \dots, n$ and $i, j = 1, 2$. Together with (2.6), we consider an initial problem

$$y(k) = \varphi^*(k), \quad (2.7)$$

$k \in \mathbb{Z}_{-m_n}^0$ with $\varphi^*: \mathbb{Z}_{-m_n}^0 \rightarrow \mathbb{R}^2$ where $\varphi^*(k) = S^{-1}\varphi(k)$ is the initial function corresponding to the initial function φ in (1.4).

Below, we consider all four possible cases (2.2)–(2.5) separately.

We define

$$\Phi_1(k) := (0, \varphi_1^*(k))^T, \quad \Phi_2(k) := (\varphi_2^*(k), 0)^T, \quad k \in \mathbb{Z}_{-m_n}^0. \quad (2.8)$$

Assuming that (1.1) is a weakly delayed system, by Lemma 1.4.2, the system (2.6) is weakly delayed system again.

2.1.1 The case (2.2) of two real distinct roots

In this case, we have $\Lambda = \Lambda_1$ and $\Lambda_1^k = \text{diag}(\lambda_1^k, \lambda_2^k)$. The necessary and sufficient conditions (1.11)–(1.14) for (2.6) turn into

$$b_{11}^{*l} + b_{22}^{*l} = 0, \quad (2.9)$$

$$\begin{vmatrix} b_{11}^{*l} & b_{12}^{*l} \\ b_{21}^{*l} & b_{22}^{*l} \end{vmatrix} = b_{11}^{*l}b_{22}^{*l} - b_{12}^{*l}b_{21}^{*l} = 0, \quad (2.10)$$

$$\begin{vmatrix} \lambda_1 & 0 \\ b_{21}^{*l} & b_{22}^{*l} \end{vmatrix} + \begin{vmatrix} b_{11}^{*l} & b_{12}^{*l} \\ 0 & \lambda_2 \end{vmatrix} = \lambda_1 b_{22}^{*l} + \lambda_2 b_{11}^{*l} = 0, \quad (2.11)$$

$$\begin{vmatrix} b_{11}^{*l} & b_{12}^{*l} \\ b_{21}^{*v} & b_{22}^{*v} \end{vmatrix} + \begin{vmatrix} b_{11}^{*v} & b_{12}^{*v} \\ b_{21}^{*l} & b_{22}^{*l} \end{vmatrix} = 0. \quad (2.12)$$

Since $\lambda_1 \neq \lambda_2$, equations (2.9), (2.11) yield $b_{11}^{*l} = b_{22}^{*l} = 0$. Then, from (2.10), we get $b_{12}^{*l}b_{21}^{*l} = 0$, so that either $b_{21}^{*l} = 0$ or $b_{12}^{*l} = 0$. In view of assumptions $B^l \neq \Theta$, $l = 1, 2, \dots, n$ we conclude that only the following cases I, II are possible

I) $b_{11}^{*l} = b_{22}^{*l} = b_{21}^{*l} = 0$, $b_{12}^{*l} \neq 0$, $l = 1, 2, \dots, n$,

II) $b_{11}^{*l} = b_{22}^{*l} = b_{12}^{*l} = 0$, $b_{21}^{*l} \neq 0$, $l = 1, 2, \dots, n$.

In Theorem 1.5 below are both cases I, II analyzed.

Theorem 2.1.1. *Let (1.1) be a weakly delayed system and equation (2.1) has two real distinct roots λ_1, λ_2 . If the case I) hold, then the solution of the initial problem (1.1), (1.4) is $x(k) = Sy(k)$, $k \in \mathbb{Z}_{-m_n}^\infty$ where $y(k)$ has the form*

$$y(k) = \begin{cases} \varphi^*(k) & \text{if } k \in \mathbb{Z}_{-m_n}^0, \\ \Lambda_1^k \varphi^*(0) + \sum_{r=0}^{k-1} \lambda_1^{k-1-r} \left[\sum_{l=1}^n b_{12}^{*l} \Phi_2(r - m_l) \right] & \text{if } k \in \mathbb{Z}_1^{m_1+1}, \\ \dots \\ \Lambda_1^k \varphi^*(0) + \sum_{r=0}^{k-1} \lambda_1^{k-1-r} \left[\sum_{l=s+1}^n b_{12}^{*l} \Phi_2(r - m_l) \right] \\ + \sum_{l=1}^s b_{12}^{*l} \left[\sum_{r=0}^{m_l} \lambda_1^{k-1-r} \Phi_2(r - m_l) \right. \\ \left. + \Phi_2(0) \sum_{r=m_l+1}^{k-1} \lambda_1^{k-1-r} \lambda_2^{r-m_l} \right] \\ \text{if } k \in \mathbb{Z}_{m_s+2}^{m_{s+1}+1}, s = 1, 2, \dots, n-1, \\ \dots \\ \Lambda_1^k \varphi^*(0) + \sum_{l=1}^n b_{12}^{*l} \left[\sum_{r=0}^{m_l} \lambda_1^{k-1-r} \Phi_2(r - m_l) \right. \\ \left. + \Phi_2(0) \sum_{r=m_l+1}^{k-1} \lambda_1^{k-1-r} \lambda_2^{r-m_l} \right] & \text{if } k \in \mathbb{Z}_{m_n+2}^\infty. \end{cases} \quad (2.13)$$

If the case II) is true, then the solution of initial problem (1.1), (1.4) is $x(k) =$

$Sy(k)$, $k \in \mathbb{Z}_{-m_n}^\infty$ where $y(k)$ has the form

$$y(k) = \begin{cases} \varphi^*(k) & \text{if } k \in \mathbb{Z}_{-m_n}^0, \\ \Lambda_1^k \varphi^*(0) + \sum_{r=0}^{k-1} \lambda_2^{k-1-r} \left[\sum_{l=1}^n b_{21}^{*l} \Phi_1(r - m_l) \right] & \text{if } k \in \mathbb{Z}_1^{m_1+1}, \\ \dots \\ \Lambda_1^k \varphi^*(0) + \sum_{r=0}^{k-1} \lambda_2^{k-1-r} \left[\sum_{l=s+1}^n b_{21}^{*l} \Phi_1(r - m_l) \right] \\ + \sum_{l=1}^s b_{21}^{*l} \left[\sum_{r=0}^{m_l} \lambda_2^{k-1-r} \Phi_1(r - m_l) \right. \\ \left. + \Phi_1(0) \sum_{r=m_l+1}^{k-1} \lambda_1^{r-m_l} \lambda_2^{k-1-r} \right] & \text{if } k \in \mathbb{Z}_{m_s+2}^{m_s+1+1}, s = 1, 2, \dots, n-1, \\ \dots \\ \Lambda_1^k \varphi^*(0) + \sum_{l=1}^n b_{21}^{*l} \left[\sum_{r=0}^{m_l} \lambda_2^{k-1-r} \Phi_1(r - m_l) \right. \\ \left. + \Phi_1(0) \sum_{r=m_l+1}^{k-1} \lambda_1^{r-m_l} \lambda_2^{k-1-r} \right] & \text{if } k \in \mathbb{Z}_{m_n+2}^\infty. \end{cases} \quad (2.14)$$

Proof. If the case I) is true, then the transformed system (2.6) takes the form

$$y_1(k+1) = \lambda_1 y_1(k) + \sum_{l=1}^n b_{12}^{*l} y_2(k - m_l), \quad (2.15)$$

$$y_2(k+1) = \lambda_2 y_2(k), \quad (2.16)$$

$$k \in \mathbb{Z}_0^\infty,$$

and if the case II) holds, then (2.6) takes the form

$$y_1(k+1) = \lambda_1 y_1(k), \quad (2.17)$$

$$y_2(k+1) = \lambda_2 y_2(k) + \sum_{l=1}^n b_{21}^{*l} y_1(k - m_l), \quad (2.18)$$

$$k \in \mathbb{Z}_0^\infty.$$

We investigate only the initial problem (2.15), (2.16), (2.7) since the initial problem (2.17), (2.18), (2.7) can be examined in a similar way.

From (2.16), (2.7), we get

$$y_2(k) = \begin{cases} \varphi_2^*(k) & \text{if } k \in \mathbb{Z}_{-m_n}^0, \\ \lambda_2^k \varphi_2^*(0) & \text{if } k \in \mathbb{Z}_1^\infty. \end{cases} \quad (2.19)$$

Then (2.15) becomes

$$y_1(k+1) = \begin{cases} \lambda_1 y_1(k) + \sum_{l=1}^n b_{12}^{*l} \varphi_2^*(k - m_l) \\ \quad \text{if } k \in \mathbb{Z}_0^{m_1}, \\ \lambda_1 y_1(k) + b_{12}^{*1} \lambda_2^{k-m_1} \varphi_2^*(0) + \sum_{l=2}^n b_{12}^{*l} \varphi_2^*(k - m_l) \\ \quad \text{if } k \in \mathbb{Z}_{m_1+1}^{m_2}, \\ \lambda_1 y_1(k) + \sum_{l=1}^2 b_{12}^{*l} \lambda_2^{k-m_l} \varphi_2^*(0) + \sum_{l=3}^n b_{12}^{*l} \varphi_2^*(k - m_l) \\ \quad \text{if } k \in \mathbb{Z}_{m_2+1}^{m_3}, \\ \dots \\ \lambda_1 y_1(k) + \sum_{l=1}^s b_{12}^{*l} \lambda_2^{k-m_l} \varphi_2^*(0) + \sum_{l=s+1}^n b_{12}^{*l} \varphi_2^*(k - m_l) \\ \quad \text{if } k \in \mathbb{Z}_{m_{s+1}}^{m_{s+1}}, s = 3, 4, \dots, n-1, \\ \dots \\ \lambda_1 y_1(k) + \sum_{l=1}^n b_{12}^{*l} \lambda_2^{k-m_l} \varphi_2^*(0) \\ \quad \text{if } k \in \mathbb{Z}_{m_n+1}^{\infty}. \end{cases} \quad (2.20)$$

First, we solve this equation for $k \in \mathbb{Z}_0^{m_1}$. This means that we consider the problem

$$\begin{cases} y_1(k+1) = \lambda_1 y_1(k) + \sum_{l=1}^n b_{12}^{*l} \varphi_2^*(k - m_l), & k \in \mathbb{Z}_0^{m_1}, \\ y_1(0) = \varphi_1^*(0). \end{cases}$$

With the aid of formula (1.20), we get

$$y_1(k) = \lambda_1^k \varphi_1^*(0) + \sum_{r=0}^{k-1} \lambda_1^{k-1-r} \left[\sum_{l=1}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right], \quad k \in \mathbb{Z}_1^{m_1+1}. \quad (2.21)$$

Now we solve equation (2.20) for $k \in \mathbb{Z}_{m_1+1}^{m_2}$ with initial data deduced from (2.21), i.e., we consider the problem.

$$\begin{cases} y_1(k+1) = \lambda_1 y_1(k) + b_{12}^{*1} \lambda_2^{k-m_1} \varphi_2^*(0) + \sum_{l=2}^n b_{12}^{*l} \varphi_2^*(k - m_l), & k \in \mathbb{Z}_{m_1+1}^{m_2}, \\ y_1(m_1+1) = \lambda_1^{m_1+1} \varphi_1^*(0) + \sum_{r=0}^{m_1} \lambda_1^{m_1-r} \left[\sum_{l=1}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right]. \end{cases}$$

Applying formula (1.20) we get (for $k \in \mathbb{Z}_{m_1+2}^{m_2+1}$)

$$\begin{aligned} y_1(k) &= \lambda_1^{k-(m_1+1)} y_1(m_1+1) \\ &+ \sum_{r=m_1+1}^{k-1} \lambda_1^{k-1-r} \left[b_{12}^{*1} \lambda_2^{r-m_1} \varphi_2^*(0) + \sum_{l=2}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right] \end{aligned}$$

$$\begin{aligned}
 &= \lambda_1^{k-m_1-1} \left[\lambda_1^{m_1+1} \varphi_1^*(0) + \sum_{r=0}^{m_1} \lambda_1^{m_1-r} \left[\sum_{l=1}^n b_{12}^{*l} \varphi_2^*(r-m_l) \right] \right] \\
 &\quad + \sum_{r=m_1+1}^{k-1} \lambda_1^{k-1-r} \left[b_{12}^{*1} \lambda_2^{r-m_1} \varphi_2^*(0) + \sum_{l=2}^n b_{12}^{*l} \varphi_2^*(r-m_l) \right] \\
 &= \lambda_1^k \varphi_1^*(0) + \sum_{r=0}^{m_1} \lambda_1^{k-1-r} \left[\sum_{l=1}^n b_{12}^{*l} \varphi_2^*(r-m_l) \right] \\
 &\quad + \sum_{r=m_1+1}^{k-1} \lambda_1^{k-1-r} \left[b_{12}^{*1} \lambda_2^{r-m_1} \varphi_2^*(0) + \sum_{l=2}^n b_{12}^{*l} \varphi_2^*(r-m_l) \right] \\
 &= \lambda_1^k \varphi_1^*(0) + \sum_{r=0}^{k-1} \lambda_1^{k-1-r} \left[\sum_{l=2}^n b_{12}^{*l} \varphi_2^*(r-m_l) \right] \\
 &\quad + b_{12}^{*1} \left[\sum_{r=0}^{m_1} \lambda_1^{k-1-r} \varphi_2^*(r-m_1) + \varphi_2^*(0) \sum_{r=m_1+1}^{k-1} \lambda_1^{k-1-r} \lambda_2^{r-m_1} \right]. \tag{2.22}
 \end{aligned}$$

Now we solve equation (2.20) for $k \in \mathbb{Z}_{m_2+1}^{m_3}$ with initial data deduced from (2.22), i.e., we consider the problem.

$$\begin{cases}
 y_1(k+1) = \lambda_1 y_1(k) + \sum_{l=1}^2 b_{12}^{*l} \lambda_2^{k-m_l} \varphi_2^*(0) + \sum_{l=3}^n b_{12}^{*l} \varphi_2^*(k-m_l), & k \in \mathbb{Z}_{m_2+1}^{m_3}, \\
 y_1(m_2+1) = \lambda_1^{m_2+1} \varphi_1^*(0) + \sum_{r=0}^{m_2} \lambda_1^{m_2-r} \left[\sum_{l=2}^n b_{12}^{*l} \varphi_2^*(r-m_l) \right] \\
 \quad + b_{12}^{*1} \left[\sum_{r=0}^{m_1} \lambda_1^{m_2-r} \varphi_2^*(r-m_1) + \varphi_2^*(0) \sum_{r=m_1+1}^{m_2} \lambda_1^{m_2-r} \lambda_2^{r-m_1} \right].
 \end{cases}$$

Applying formula (1.20) yields (for $k \in \mathbb{Z}_{m_2+2}^{m_3+1}$)

$$\begin{aligned}
 y_1(k) &= \lambda_1^{k-(m_2+1)} y_1(m_2+1) \\
 &\quad + \sum_{r=m_2+1}^{k-1} \lambda_1^{k-1-r} \left[\sum_{l=1}^2 b_{12}^{*l} \lambda_2^{r-m_l} \varphi_2^*(0) + \sum_{l=3}^n b_{12}^{*l} \varphi_2^*(r-m_l) \right] \\
 &= \lambda_1^{k-m_2-1} \left[\lambda_1^{m_2+1} \varphi_1^*(0) + \sum_{r=0}^{m_2} \lambda_1^{m_2-r} \left[\sum_{l=2}^n b_{12}^{*l} \varphi_2^*(r-m_l) \right] \right] \\
 &\quad + b_{12}^{*1} \left[\sum_{r=0}^{m_1} \lambda_1^{m_2-r} \varphi_2^*(r-m_1) + \varphi_2^*(0) \sum_{r=m_1+1}^{m_2} \lambda_1^{m_2-r} \lambda_2^{r-m_1} \right] \\
 &\quad + \sum_{r=m_2+1}^{k-1} \lambda_1^{k-1-r} \left[\sum_{l=1}^2 b_{12}^{*l} \lambda_2^{r-m_l} \varphi_2^*(0) + \sum_{l=3}^n b_{12}^{*l} \varphi_2^*(r-m_l) \right] \\
 &= \lambda_1^k \varphi_1^*(0) + \sum_{r=0}^{m_2} \lambda_1^{k-1-r} \left[\sum_{l=2}^n b_{12}^{*l} \varphi_2^*(r-m_l) \right] \\
 &\quad + b_{12}^{*1} \left[\sum_{r=0}^{m_1} \lambda_1^{k-1-r} \varphi_2^*(r-m_1) + \varphi_2^*(0) \sum_{r=m_1+1}^{m_2} \lambda_1^{k-1-r} \lambda_2^{r-m_1} \right] \\
 &\quad + \sum_{r=m_2+1}^{k-1} \lambda_1^{k-1-r} \left[\sum_{l=1}^2 b_{12}^{*l} \lambda_2^{r-m_l} \varphi_2^*(0) + \sum_{l=3}^n b_{12}^{*l} \varphi_2^*(r-m_l) \right]
 \end{aligned}$$

$$\begin{aligned}
 &= \lambda_1^k \varphi_1^*(0) + \sum_{r=0}^{k-1} \lambda_1^{k-1-r} \left[\sum_{l=3}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right] \\
 &\quad + b_{12}^{*1} \left[\sum_{r=0}^{m_1} \lambda_1^{k-1-r} \varphi_2^*(r - m_1) + \varphi_2^*(0) \sum_{r=m_1+1}^{k-1} \lambda_1^{k-1-r} \lambda_2^{r-m_1} \right] \\
 &\quad + b_{12}^{*2} \left[\sum_{r=0}^{m_2} \lambda_1^{k-1-r} \varphi_2^*(r - m_2) + \varphi_2^*(0) \sum_{r=m_2+1}^{k-1} \lambda_1^{k-1-r} \lambda_2^{r-m_2} \right]. \tag{2.23}
 \end{aligned}$$

From (2.21), (2.22) and (2.23) we deduce that expected form of the solution of the initial problem for $k \in \mathbb{Z}_{m_{s-1}+1}^{m_s}$ with initial data derived from the solution of previous equation for $k \in \mathbb{Z}_{m_{s-2}+1}^{m_{s-1}}$ is

$$\begin{aligned}
 y_1(k) &= \lambda_1^k \varphi_1^*(0) + \sum_{r=0}^{k-1} \lambda_1^{k-1-r} \left[\sum_{l=s}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right] \tag{2.24} \\
 &\quad + \sum_{l=1}^{s-1} b_{12}^{*l} \left[\sum_{r=0}^{m_l} \lambda_1^{k-1-r} \varphi_2^*(r - m_l) + \varphi_2^*(0) \sum_{r=m_l+1}^{k-1} \lambda_1^{k-1-r} \lambda_2^{r-m_l} \right] \\
 &\quad \text{if } k \in \mathbb{Z}_{m_{s-1}+2}^{m_s+1}.
 \end{aligned}$$

We solve equation (2.20) for $k \in \mathbb{Z}_{m_s+1}^{m_s+1}$ with initial data deduced from (2.24), i.e., we consider the problem

$$\left\{ \begin{aligned}
 y_1(k+1) &= \lambda_1 y_1(k) + \sum_{l=1}^s b_{12}^{*l} \lambda_2^{k-m_l} \varphi_2^*(0) + \sum_{l=s+1}^n b_{12}^{*l} \varphi_2^*(k - m_l), \quad k \in \mathbb{Z}_{m_s+1}^{m_s+1}, \\
 y_1(m_s+1) &= \lambda_1^{m_s+1} \varphi_1^*(0) + \sum_{r=0}^{m_s} \lambda_1^{m_s-r} \left[\sum_{l=s}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right] \\
 &\quad + \sum_{l=1}^{s-1} b_{12}^{*l} \left[\sum_{r=0}^{m_l} \lambda_1^{m_s-r} \varphi_2^*(r - m_l) + \varphi_2^*(0) \sum_{r=m_l+1}^{m_s} \lambda_1^{m_s-r} \lambda_2^{r-m_l} \right].
 \end{aligned} \right.$$

Applying formula (1.20) yields (for $k \in \mathbb{Z}_{m_s+2}^{m_s+1+1}$)

$$\begin{aligned}
 y_1(k) &= \lambda_1^{k-(m_s+1)} y_1(m_s+1) \\
 &\quad + \sum_{r=m_s+1}^{k-1} \lambda_1^{k-1-r} \left[\sum_{l=1}^s b_{12}^{*l} \lambda_2^{r-m_l} \varphi_2^*(0) + \sum_{l=s+1}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right] \\
 &= \lambda_1^{k-(m_s+1)} \left[\lambda_1^{m_s+1} \varphi_1^*(0) + \sum_{r=0}^{m_s} \lambda_1^{m_s-r} \left[\sum_{l=s}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right] \right] \\
 &\quad + \sum_{l=1}^{s-1} b_{12}^{*l} \left[\sum_{r=0}^{m_l} \lambda_1^{m_s-r} \varphi_2^*(r - m_l) + \varphi_2^*(0) \sum_{r=m_l+1}^{m_s} \lambda_1^{m_s-r} \lambda_2^{r-m_l} \right] \\
 &\quad + \sum_{r=m_s+1}^{k-1} \lambda_1^{k-1-r} \left[\sum_{l=1}^s b_{12}^{*l} \lambda_2^{r-m_l} \varphi_2^*(0) + \sum_{l=s+1}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right] \\
 &= \lambda_1^k \varphi_1^*(0) + \sum_{r=0}^{m_s} \lambda_1^{k-1-r} \left[\sum_{l=s}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right]
 \end{aligned}$$

$$\begin{aligned}
 & + \sum_{l=1}^{s-1} b_{12}^{*l} \left[\sum_{r=0}^{m_l} \lambda_1^{k-1-r} \varphi_2^*(r - m_l) + \varphi_2^*(0) \sum_{r=m_l+1}^{m_s} \lambda_1^{k-1-r} \lambda_2^{r-m_l} \right] \\
 & + \sum_{r=m_s+1}^{k-1} \lambda_1^{k-1-r} \left[\sum_{l=1}^s b_{12}^{*l} \lambda_2^{r-m_l} \varphi_2^*(0) + \sum_{l=s+1}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right] \\
 = & \lambda_1^k \varphi_1^*(0) + \sum_{r=0}^{m_s} \lambda_1^{k-1-r} \left[b_{12}^{*s} \varphi_2^*(r - m_s) + \sum_{l=s+1}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right] \\
 & + b_{12}^{*1} \left[\sum_{r=0}^{m_1} \lambda_1^{k-1-r} \varphi_2^*(r - m_1) + \varphi_2^*(0) \sum_{r=m_1+1}^{m_s} \lambda_1^{k-1-r} \lambda_2^{r-m_1} \right] \\
 & + b_{12}^{*2} \left[\sum_{r=0}^{m_2} \lambda_1^{k-1-r} \varphi_2^*(r - m_2) + \varphi_2^*(0) \sum_{r=m_2+1}^{m_s} \lambda_1^{k-1-r} \lambda_2^{r-m_2} \right] \\
 & + \dots \\
 & + b_{12}^{*s-1} \left[\sum_{r=0}^{m_{s-1}} \lambda_1^{k-1-r} \varphi_2^*(r - m_{s-1}) + \varphi_2^*(0) \sum_{r=m_{s-1}+1}^{m_s} \lambda_1^{k-1-r} \lambda_2^{r-m_{s-1}} \right] \\
 & + b_{12}^{*1} \left[\varphi_2^*(0) \sum_{r=m_s+1}^{k-1} \lambda_1^{k-1-r} \lambda_2^{r-m_1} \right] + b_{12}^{*2} \left[\varphi_2^*(0) \sum_{r=m_s+1}^{k-1} \lambda_1^{k-1-r} \lambda_2^{r-m_2} \right] \\
 & + \dots \\
 & + b_{12}^{*s} \left[\varphi_2^*(0) \sum_{r=m_s+1}^{k-1} \lambda_1^{k-1-r} \lambda_2^{r-m_s} \right] + \sum_{r=m_s+1}^{k-1} \lambda_1^{k-1-r} \left[\sum_{l=s+1}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right] \\
 = & \lambda_1^k \varphi_1^*(0) + \sum_{r=0}^{k-1} \lambda_1^{k-1-r} \left[\sum_{l=s+1}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right] \\
 & + \sum_{l=1}^s b_{12}^{*l} \left[\sum_{r=0}^{m_l} \lambda_1^{k-1-r} \varphi_2^*(r - m_l) + \varphi_2^*(0) \sum_{r=m_l+1}^{m_s} \lambda_1^{k-1-r} \lambda_2^{r-m_l} \right]. \tag{2.25}
 \end{aligned}$$

In the end we solve equation (2.20) for $k \in \mathbb{Z}_{m_n+1}^\infty$ with initial data deduced from (2.25), i.e., we consider the problem

$$\begin{cases}
 y_1(k+1) = \lambda_1 y_1(k) + \sum_{l=1}^n b_{12}^{*l} \lambda_2^{k-m_l} \varphi_2^*(0), & k \in \mathbb{Z}_{m_n+1}^\infty, \\
 y_1(m_n+1) = \lambda_1^{m_n+1} \varphi_1^*(0) + \sum_{r=0}^{m_n} \lambda_1^{m_n-r} b_{12}^{*n} \varphi_2^*(r - m_n) \\
 \quad + \sum_{l=1}^{n-1} b_{12}^{*l} \left[\sum_{r=0}^{m_l} \lambda_1^{m_n-r} \varphi_2^*(r - m_l) + \varphi_2^*(0) \sum_{r=m_l+1}^{m_n} \lambda_1^{m_n-r} \lambda_2^{r-m_l} \right].
 \end{cases}$$

Applying formula (1.20) yields (for $k \in \mathbb{Z}_{m_n+2}^\infty$)

$$\begin{aligned}
 y_1(k) & = \lambda_1^{k-(m_n+1)} y_1(m_n+1) + \sum_{r=m_n+1}^{k-1} \lambda_1^{k-1-r} \left[\sum_{l=1}^n b_{12}^{*l} \lambda_2^{r-m_l} \varphi_2^*(0) \right] \\
 & = \lambda_1^{k-(m_n+1)} \left[\lambda_1^{m_n+1} \varphi_1^*(0) + \sum_{r=0}^{m_n} \lambda_1^{m_n-r} b_{12}^{*n} \varphi_2^*(r - m_n) \right. \\
 & \quad \left. + \sum_{l=1}^{n-1} b_{12}^{*l} \left[\sum_{r=0}^{m_l} \lambda_1^{m_n-r} \varphi_2^*(r - m_l) + \varphi_2^*(0) \sum_{r=m_l+1}^{m_n} \lambda_1^{m_n-r} \lambda_2^{r-m_l} \right] \right]
 \end{aligned}$$

$$\begin{aligned}
 & + \sum_{r=m_n+1}^{k-1} \lambda_1^{k-1-r} \left[\sum_{l=1}^n b_{12}^{*l} \lambda_2^{r-m_l} \varphi_2^*(0) \right] \\
 = & \lambda_1^k \varphi_1^*(0) + \sum_{r=0}^{m_n} \lambda_1^{k-1-r} b_{12}^{*n} \varphi_2^*(r - m_n) \\
 & + \sum_{l=1}^{n-1} b_{12}^{*l} \left[\sum_{r=0}^{m_l} \lambda_1^{k-1-r} \varphi_2^*(r - m_l) + \varphi_2^*(0) \sum_{r=m_l+1}^{m_n} \lambda_1^{k-1-r} \lambda_2^{r-m_l} \right] \\
 & + \sum_{r=m_n+1}^{k-1} \lambda_1^{k-1-r} \left[\sum_{l=1}^n b_{12}^{*l} \lambda_2^{r-m_l} \varphi_2^*(0) \right] \\
 = & \lambda_1^k \varphi_1^*(0) + \sum_{r=0}^{m_n} \lambda_1^{k-1-r} b_{12}^{*n} \varphi_2^*(r - m_n) \\
 & + b_{12}^{*1} \left[\sum_{r=0}^{m_1} \lambda_1^{k-1-r} \varphi_2^*(r - m_1) + \varphi_2^*(0) \sum_{r=m_1+1}^{m_s} \lambda_1^{k-1-r} \lambda_2^{r-m_1} \right] \\
 & + b_{12}^{*2} \left[\sum_{r=0}^{m_2} \lambda_1^{k-1-r} \varphi_2^*(r - m_2) + \varphi_2^*(0) \sum_{r=m_2+1}^{m_s} \lambda_1^{k-1-r} \lambda_2^{r-m_2} \right] \\
 & + \dots \\
 & + b_{12}^{*n-1} \left[\sum_{r=0}^{m_{n-1}} \lambda_1^{k-1-r} \varphi_2^*(r - m_{n-1}) + \varphi_2^*(0) \sum_{r=m_{n-1}+1}^{m_s} \lambda_1^{k-1-r} \lambda_2^{r-m_{n-1}} \right] \\
 & + b_{12}^{*1} \left[\varphi_2^*(0) \sum_{r=m_s+1}^{k-1} \lambda_1^{k-1-r} \lambda_2^{r-m_1} \right] + b_{12}^{*2} \left[\varphi_2^*(0) \sum_{r=m_s+1}^{k-1} \lambda_1^{k-1-r} \lambda_2^{r-m_2} \right] \\
 & + \dots \\
 & + b_{12}^{*n} \left[\varphi_2^*(0) \sum_{r=m_n+1}^{k-1} \lambda_1^{k-1-r} \lambda_2^{r-m_n} \right] \\
 = & \lambda_1^k \varphi_1^*(0) + \sum_{l=1}^n b_{12}^{*l} \left[\sum_{r=0}^{m_l} \lambda_1^{k-1-r} \varphi_2^*(r - m_l) + \varphi_2^*(0) \sum_{r=m_l+1}^{k-1} \lambda_1^{k-1-r} \lambda_2^{r-m_l} \right]. \quad (2.26)
 \end{aligned}$$

Summing up all particular cases (2.21)–(2.26) we have

$$y_1(k) = \left\{ \begin{array}{l}
 \varphi_1^*(k) \quad \text{if } k \in \mathbb{Z}_{-m_n}^0, \\
 \lambda_1^k \varphi_1^*(0) + \sum_{r=0}^{k-1} \lambda_1^{k-1-r} \left[\sum_{l=1}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right] \\
 \quad \text{if } k \in \mathbb{Z}_1^{m_1+1}, \\
 \lambda_1^k \varphi_1^*(0) + \sum_{r=0}^{k-1} \lambda_1^{k-1-r} \left[\sum_{l=2}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right] \\
 + b_{12}^{*1} \left[\sum_{r=0}^{m_1} \lambda_1^{k-1-r} \varphi_2^*(r - m_1) + \varphi_2^*(0) \sum_{r=m_1+1}^{k-1} \lambda_1^{k-1-r} \lambda_2^{r-m_1} \right] \\
 \quad \text{if } k \in \mathbb{Z}_{m_1+2}^{m_2+1}, \\
 \lambda_1^k \varphi_1^*(0) + \sum_{r=0}^{k-1} \lambda_1^{k-1-r} \left[\sum_{l=3}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right] \\
 + b_{12}^{*1} \left[\sum_{r=0}^{m_1} \lambda_1^{k-1-r} \varphi_2^*(r - m_1) + \varphi_2^*(0) \sum_{r=m_1+1}^{k-1} \lambda_1^{k-1-r} \lambda_2^{r-m_1} \right] \\
 + b_{12}^{*2} \left[\sum_{r=0}^{m_2} \lambda_1^{k-1-r} \varphi_2^*(r - m_2) + \varphi_2^*(0) \sum_{r=m_2+1}^{k-1} \lambda_1^{k-1-r} \lambda_2^{r-m_2} \right] \\
 \quad \text{if } k \in \mathbb{Z}_{m_2+2}^{m_3+1}, \\
 \dots \\
 \lambda_1^k \varphi_1^*(0) + \sum_{r=0}^{k-1} \lambda_1^{k-1-r} \left[\sum_{l=s+1}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right] \\
 + \sum_{l=1}^s b_{12}^{*l} \left[\sum_{r=0}^{m_l} \lambda_1^{k-1-r} \varphi_2^*(r - m_l) \right. \\
 \quad \left. + \varphi_2^*(0) \sum_{r=m_l+1}^{k-1} \lambda_1^{k-1-r} \lambda_2^{r-m_l} \right] \\
 \quad \text{if } k \in \mathbb{Z}_{m_s+2}^{m_{s+1}+1}, \\
 \dots \\
 \lambda_1^k \varphi_1^*(0) + \sum_{l=1}^n b_{12}^{*l} \left[\sum_{r=0}^{m_l} \lambda_1^{k-1-r} \varphi_2^*(r - m_l) \right. \\
 \quad \left. + \varphi_2^*(0) \sum_{r=m_l+1}^{k-1} \lambda_1^{k-1-r} \lambda_2^{r-m_l} \right] \\
 \quad \text{if } k \in \mathbb{Z}_{m_n+2}^\infty.
 \end{array} \right. \quad (2.27)$$

Now, taking into account (2.8), formula (2.13) is a consequence of (2.19) and (2.27). Formula (2.14) can be proved in a similar way.

Finally, we note that both formulas (2.13), (2.14) remain valid for $b_{12}^{*l} = b_{21}^{*l} = 0$. In this case, the transformed system (1.1) reduces to a system without delays. This possibility is excluded by conditions (1.3).

□

2.1.2 The case (2.3) of two complex conjugate roots

The necessary and sufficient conditions (1.11)–(1.14) take the forms (2.9), (2.10), (2.12) and

$$\begin{vmatrix} p & q \\ b_{21}^{*l} & b_{22}^{*l} \end{vmatrix} + \begin{vmatrix} b_{11}^{*l} & b_{12}^{*l} \\ -q & p \end{vmatrix} = p(b_{11}^{*l} + b_{22}^{*l}) + q(b_{12}^{*l} - b_{21}^{*l}) = 0 \quad (2.28)$$

where $l, v = 1, 2, \dots, n$ and $v > l$.

The system of conditions (2.9), (2.10) and (2.28) gives $b_{12}^{*l} = b_{21}^{*l}$, $(b_{11}^{*l})^2 = -(b_{12}^{*l})^2$ and admits only one possibility, namely,

$$b_{11}^{*l} = b_{22}^{*l} = b_{12}^{*l} = b_{21}^{*l} = 0.$$

Consequently, $B^{*l} = \Theta$, $B^l = \Theta$.

The initial problem (1.1), (1.4) reduces to a problem without delay

$$\begin{cases} x(k+1) = Ax(k), \\ x(k) = \varphi(k), \quad k \in \mathbb{Z}_{-m_n}^0 \end{cases}$$

and, obviously,

$$x(k) = \begin{cases} \varphi(k) & \text{if } k \in \mathbb{Z}_{-m_n}^0, \\ A^k \varphi(0) & \text{if } k \in \mathbb{Z}_1^\infty. \end{cases} \quad (2.29)$$

From this discussion, the next theorem follows.

Theorem 2.1.2. *There exists no weakly delayed system (1.1) if Λ has the form (2.3).*

Finally, we note that the assumptions (1.3) alone exclude this case.

2.1.3 The case (2.4) of double real root

In this case we have $\Lambda = \Lambda_3$ and $\Lambda_3^k = \text{diag}(\lambda^k, \lambda^k)$. For (2.6), the necessary and sufficient conditions (1.11)–(1.14) are reduced to (2.9), (2.10), (2.12) and

$$\begin{vmatrix} \lambda & 0 \\ b_{21}^{*l} & b_{22}^{*l} \end{vmatrix} + \begin{vmatrix} b_{11}^{*l} & b_{12}^{*l} \\ 0 & \lambda \end{vmatrix} = \lambda(b_{11}^{*l} + b_{22}^{*l}) = 0 \quad (2.30)$$

where $l = 1, 2, \dots, n$.

From (2.9), (2.10) and (2.30), we get $b_{12}^{*l} b_{21}^{*l} = -(b_{11}^{*l})^2$. From the condition (2.12) we get

$$b_{11}^{*l} b_{22}^{*v} - b_{12}^{*l} b_{21}^{*v} + b_{22}^{*l} b_{11}^{*v} - b_{21}^{*l} b_{12}^{*v} = 0 \quad (2.31)$$

where $l, v = 1, 2, \dots, n$ and $v > l$. Multiplying (2.31) by $b_{12}^{*l}b_{12}^{*v}$, we have

$$b_{11}^{*l}b_{22}^{*v}b_{12}^{*l}b_{12}^{*v} - (b_{12}^{*l})^2b_{21}^{*v}b_{12}^{*v} + b_{22}^{*l}b_{11}^{*v}b_{12}^{*l}b_{12}^{*v} - b_{21}^{*l}b_{12}^{*l}(b_{12}^{*v})^2 = 0. \quad (2.32)$$

Substituting $b_{12}^{*l}b_{21}^{*l} = -(b_{11}^{*l})^2$, $b_{12}^{*v}b_{21}^{*v} = -(b_{11}^{*v})^2$ into (2.32) and using (2.9) we obtain

$$-b_{11}^{*l}b_{11}^{*v}b_{12}^{*l}b_{12}^{*v} + (b_{12}^{*l})^2(b_{11}^{*v})^2 - b_{11}^{*l}b_{11}^{*v}b_{12}^{*l}b_{12}^{*v} + (b_{11}^{*l})^2(b_{12}^{*v})^2 = 0. \quad (2.33)$$

The equation (2.33) can be written as

$$(b_{12}^{*l}b_{11}^{*v} - b_{12}^{*v}b_{11}^{*l})^2 = 0$$

and

$$b_{12}^{*l}b_{11}^{*v} = b_{12}^{*v}b_{11}^{*l}. \quad (2.34)$$

We analyse the two possible cases: $b_{12}^{*l}b_{21}^{*l} = 0$ and $b_{12}^{*l}b_{21}^{*l} \neq 0$.

For the case $b_{12}^{*l}b_{21}^{*l} = 0$, we have from (2.9), (2.10) that $b_{11}^{*l} = b_{22}^{*l} = 0$ and $b_{12}^{*l} = 0$ or $b_{21}^{*l} = 0$. For $b_{12}^{*l} = 0$ and $b_{21}^{*l} \neq 0$, condition (2.12) gives $b_{12}^{*v} = 0$, where $l, v = 1, 2, \dots, n$ and $v > l$. Then, from (2.9), (2.10) for $l = v$, we get $b_{11}^{*v} = b_{22}^{*v} = 0$ and $b_{21}^{*v} \neq 0$.

For $b_{21}^{*l} = 0$ and $b_{12}^{*l} \neq 0$, condition (2.12) gives $b_{21}^{*v} = 0$, where $l, v = 1, 2, \dots, n$ and $v > l$. Then, from (2.9), (2.10) for $l = v$, we get $b_{11}^{*v} = b_{22}^{*v} = 0$ and $b_{12}^{*v} \neq 0$.

Now we discuss the case $b_{12}^{*l}b_{21}^{*l} \neq 0$. From conditions (2.9), (2.10), we have $b_{12}^{*l}b_{21}^{*l} = -(b_{11}^{*l})^2$ and $b_{11}^{*l}b_{22}^{*l} \neq 0$. This yields $b_{11}^{*l} \neq 0$, $b_{22}^{*l} \neq 0$ and, from (2.34), we have $b_{11}^{*v} \neq 0$, $b_{12}^{*v} \neq 0$. By conditions (2.9), (2.10) for $v = l$, we get $b_{22}^{*v} \neq 0$, $b_{21}^{*v} \neq 0$.

From the assumptions $B^l \neq \Theta$, we conclude that only the following cases I, II, III are possible

$$\text{I)} \quad b_{11}^{*l} = b_{22}^{*l} = b_{21}^{*l} = 0, \quad b_{12}^{*l} \neq 0,$$

$$\text{II)} \quad b_{11}^{*l} = b_{22}^{*l} = b_{12}^{*l} = 0, \quad b_{21}^{*l} \neq 0,$$

$$\text{III)} \quad b_{12}^{*l}b_{21}^{*l} \neq 0,$$

where $l = 1, 2, \dots, n$.

The case $b_{12}^{*l}b_{21}^{*l} = 0$.

Theorem 2.1.3. *Let (1.1) be a weakly delayed system, equation (2.1) has a two-fold root $\lambda_{1,2} = \lambda$, $b_{12}^{*l}b_{21}^{*l} = 0$ and the matrix Λ has the form (2.4). Then the solution of*

the initial problem (1.1), (1.4) is $x(k) = Sy(k)$, $k \in \mathbb{Z}_{-m_n}^\infty$ where in the case $b_{21}^{*l} = 0$, $y(k)$ has the form

$$y(k) = \begin{cases} \varphi^*(k) & \text{if } k \in \mathbb{Z}_{-m_n}^0, \\ \Lambda_3^k \varphi^*(0) + \sum_{r=0}^{k-1} \lambda^{k-1-r} \left[\sum_{l=1}^n b_{12}^{*l} \Phi_2(r - m_l) \right] & \text{if } k \in \mathbb{Z}_1^{m_1+1}, \\ \dots \\ \Lambda_3^k \varphi^*(0) + \sum_{r=0}^{k-1} \lambda^{k-1-r} \left[\sum_{l=s+1}^n b_{12}^{*l} \Phi_2(r - m_l) \right. \\ \quad \left. + \sum_{l=1}^s b_{12}^{*l} \left[\sum_{r=0}^{m_l} \lambda^{k-1-r} \Phi_2(r - m_l) \right. \right. \\ \quad \quad \left. \left. + (k-1-m_l) \lambda^{k-1-m_l} \Phi_2(0) \right] \right] & \text{if } k \in \mathbb{Z}_{m_s+2}^{m_{s+1}+1}, \quad s = 1, 2, \dots, n-1, \\ \dots \\ \Lambda_3^k \varphi^*(0) + \sum_{l=1}^n b_{12}^{*l} \left[\sum_{r=0}^{m_l} \lambda^{k-1-r} \Phi_2(r - m_l) \right. \\ \quad \left. + (k-1-m_l) \lambda^{k-1-m_l} \Phi_2(0) \right] & \text{if } k \in \mathbb{Z}_{m_n+2}^\infty. \end{cases} \quad (2.35)$$

If $b_{12}^{*l} = 0$ is true then the solution of initial problem (1.1), (1.4) is $x(k) = Sy(k)$, $k \in \mathbb{Z}_{-m_n}^\infty$ where $y(k)$ has the form

$$y(k) = \begin{cases} \varphi^*(k) & \text{if } k \in \mathbb{Z}_{-m_n}^0, \\ \Lambda_3^k \varphi^*(0) + \sum_{r=0}^{k-1} \lambda^{k-1-r} \left[\sum_{l=1}^n b_{21}^{*l} \Phi_1(r - m_l) \right] & \text{if } k \in \mathbb{Z}_1^{m_1+1}, \\ \dots \\ \Lambda_3^k \varphi^*(0) + \sum_{r=0}^{k-1} \lambda^{k-1-r} \left[\sum_{l=s+1}^n b_{21}^{*l} \Phi_1(r - m_l) \right. \\ \quad \left. + \sum_{l=1}^s b_{21}^{*l} \left[\sum_{r=0}^{m_l} \lambda^{k-1-r} \Phi_1(r - m_l) \right. \right. \\ \quad \quad \left. \left. + (k-1-m_l) \lambda^{k-1-m_l} \Phi_1(0) \right] \right] & \text{if } k \in \mathbb{Z}_{m_s+2}^{m_{s+1}+1}, \quad s = 1, 2, \dots, n-1 \\ \dots \\ \Lambda_3^k \varphi^*(0) + \sum_{l=1}^n b_{21}^{*l} \left[\sum_{r=0}^{m_l} \lambda^{k-1-r} \Phi_1(r - m_l) \right. \\ \quad \left. + (k-1-m_l) \lambda^{k-1-m_l} \Phi_1(0) \right] & \text{if } k \in \mathbb{Z}_{m_n+2}^\infty. \end{cases} \quad (2.36)$$

Proof. Case I) means that $b_{12}^{*l} \neq 0$. Then (2.6) turns into the system

$$y_1(k+1) = \lambda y_1(k) + \sum_{l=1}^n b_{12}^{*l} y_2(k-m_l), \quad k \in \mathbb{Z}_0^\infty \quad (2.37)$$

$$y_2(k+1) = \lambda y_2(k), \quad (2.38)$$

and, if $b_{21}^{*l} \neq 0$, (2.6) turns into the system

$$y_1(k+1) = \lambda y_1(k), \quad (2.39)$$

$$y_2(k+1) = \lambda y_2(k) + \sum_{l=1}^n b_{21}^{*l} y_1(k-m_l), \quad k \in \mathbb{Z}_0^\infty \quad (2.40)$$

System (2.37), (2.38) can be solved in much the same way as the systems (2.15), (2.16) if we put $\lambda_1 = \lambda_2 = \lambda$, and the discussion of the system (2.39), (2.40) goes along the same lines as that of the system (2.17), (2.18) with $\lambda_1 = \lambda_2 = \lambda$. Formulas (2.35) and (2.36) are consequences of (2.13), (2.14). □

The case $b_{12}^{*l} b_{21}^{*l} \neq 0$.

For $k \in \mathbb{Z}_{-m_n}^0$, we define

$$\Phi_l^*(k) := \left(b_{11}^{*l} \left[\varphi_1^*(k) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(k) \right], -\frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(k) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(k) \right] \right)^T. \quad (2.41)$$

Theorem 2.1.4. *Let the system (1.1) be a weakly delayed system, equation (2.1) admits two repeated roots $\lambda_{1,2} = \lambda$, $b_{12}^{*l} b_{21}^{*l} \neq 0$ and the matrix Λ_3 has the form (2.4). Then the solution of the initial problem (1.1), (1.4) is given by $x(k) = Sy(k)$, $k \in$*

$\mathbb{Z}_{-m_n}^\infty$ where $y(k)$ has the form

$$y(k) = \begin{cases} \varphi^*(k) & \text{if } k \in \mathbb{Z}_{-m_n}^0, \\ \Lambda_3^k \varphi^*(0) + \sum_{r=0}^{k-1} \lambda^{k-1-r} \left[\sum_{l=1}^n \Phi_l^*(r - m_l) \right] & \text{if } k \in \mathbb{Z}_1^{m_1+1}, \\ \dots \\ \Lambda_3^k \varphi^*(0) + \sum_{r=0}^{k-1} \lambda^{k-1-r} \left[\sum_{l=s+1}^n \Phi_l^*(r - m_l) \right] \\ \quad + \sum_{l=1}^s \left[\sum_{r=0}^{m_l} \lambda^{k-1-r} \Phi_l^*(r - m_l) \right. \\ \quad \quad \left. + (k-1-m_l) \lambda^{k-1-m_l} \Phi_l^*(0) \right] & \text{if } k \in \mathbb{Z}_{m_s+2}^{m_{s+1}+1}, \quad s = 1, 2, \dots, n-1, \\ \dots \\ \Lambda_3^k \varphi^*(0) + \sum_{l=1}^n \left[\sum_{r=0}^{m_l} \lambda^{k-1-r} \Phi_l^*(r - m_l) \right. \\ \quad \left. + (k-1-m_l) \lambda^{k-1-m_l} \Phi_l^*(0) \right] & \text{if } k \in \mathbb{Z}_{m_n+2}^\infty. \end{cases} \quad (2.42)$$

Proof. In this case, all the entries of B^{*l} are nonzero and, from (2.9)–(2.10) and (2.30), we get

$$B^* = \begin{pmatrix} b_{11}^{*l} & b_{12}^{*l} \\ -(b_{11}^{*l})^2/b_{12}^{*l} & -b_{11}^{*l} \end{pmatrix}$$

where $l = 1, 2, \dots, n$. Then, the system (2.6) reduces to

$$y_1(k+1) = \lambda y_1(k) + \sum_{l=1}^n \left[b_{11}^{*l} y_1(k - m_l) + b_{12}^{*l} y_2(k - m_l) \right], \quad (2.43)$$

$$y_2(k+1) = \lambda y_2(k) - \sum_{l=1}^n \left[\frac{(b_{11}^{*l})^2}{b_{12}^{*l}} y_1(k - m_l) + b_{11}^{*l} y_2(k - m_l) \right] \quad (2.44)$$

where $k \in \mathbb{Z}_0^\infty$. It is easy to see (multiplying (2.44) by b_{12}^{*1}/b_{11}^{*1} and summing both equations) that

$$y_1(k+1) + \frac{b_{12}^{*1}}{b_{11}^{*1}} y_2(k+1) = \lambda \left[y_1(k) + \frac{b_{12}^{*1}}{b_{11}^{*1}} y_2(k) \right], \quad k \in \mathbb{Z}_0^\infty. \quad (2.45)$$

Equation (2.45) is a homogeneous equation with respect to the unknown expression

$$y_1(k) + \left(b_{12}^{*1}/b_{11}^{*1} \right) y_2(k).$$

Then, using (1.20), we obtain

$$y_1(k) + \frac{b_{12}^{*1}}{b_{11}^{*1}} y_2(k) = \begin{cases} \varphi_1^*(k) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(k) & \text{if } k \in \mathbb{Z}_{-m_n}^0, \\ \lambda^k \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] & \text{if } k \in \mathbb{Z}_1^\infty. \end{cases} \quad (2.46)$$

With the aid of (2.46), we rewrite the system (2.43), (2.44) as follows:

$$y_1(k+1) = \begin{cases} \lambda y_1(k) + \sum_{l=1}^n b_{11}^{*l} \left[\varphi_1^*(k-m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(k-m_l) \right] \\ \quad \text{if } k \in \mathbb{Z}_0^{m_1}, \\ \lambda y_1(k) + b_{11}^{*1} \lambda^{k-m_1} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \\ \quad + \sum_{l=2}^n b_{11}^{*l} \left[\varphi_1^*(k-m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(k-m_l) \right] \\ \quad \text{if } k \in \mathbb{Z}_{m_1+1}^{m_2}, \\ \lambda y_1(k) + \sum_{l=1}^2 b_{11}^{*l} \lambda^{k-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \\ \quad + \sum_{l=3}^n b_{11}^{*l} \left[\varphi_1^*(k-m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(k-m_l) \right] \\ \quad \text{if } k \in \mathbb{Z}_{m_2+1}^{m_3}, \\ \dots \\ \lambda y_1(k) + \sum_{l=1}^s b_{11}^{*l} \lambda^{k-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \\ \quad + \sum_{l=s+1}^n b_{11}^{*l} \left[\varphi_1^*(k-m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(k-m_l) \right] \\ \quad \text{if } k \in \mathbb{Z}_{m_s+1}^{m_{s+1}}, s = 3, 4, \dots, n-1, \\ \dots \\ \lambda y_1(k) + \sum_{l=1}^n b_{11}^{*l} \lambda^{k-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \\ \quad \text{if } k \in \mathbb{Z}_{m_n+1}^\infty, \end{cases} \quad (2.47)$$

$$y_2(k+1) = \left\{ \begin{array}{l} \lambda y_2(k) - \sum_{l=1}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(k - m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(k - m_l) \right] \\ \quad \text{if } k \in \mathbb{Z}_0^{m_1}, \\ \\ \lambda y_2(k) - \frac{(b_{11}^{*1})^2}{b_{12}^{*1}} \lambda^{k-m_1} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \\ \quad - \sum_{l=2}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(k - m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(k - m_l) \right] \\ \quad \text{if } k \in \mathbb{Z}_{m_1+1}^{m_2}, \\ \\ \lambda y_2(k) - \sum_{l=1}^2 \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \\ \quad - \sum_{l=3}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(k - m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(k - m_l) \right] \\ \quad \text{if } k \in \mathbb{Z}_{m_2+1}^{m_3}, \\ \\ \dots \\ \\ \lambda y_2(k) - \sum_{l=1}^s \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \\ \quad - \sum_{l=s+1}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(k - m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(k - m_l) \right] \\ \quad \text{if } k \in \mathbb{Z}_{m_{s+1}}^{m_{s+1}}, s = 3, 4, \dots, n-1, \\ \\ \dots \\ \\ \lambda y_2(k) - \sum_{l=1}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \lambda^{k-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \\ \quad \text{if } k \in \mathbb{Z}_{m_n+1}^{\infty}. \end{array} \right. \quad (2.48)$$

First, we solve this system for $k \in \mathbb{Z}_0^{m_1}$ and consider the problems

$$\left\{ \begin{array}{l} y_1(k+1) = \lambda y_1(k) + \sum_{l=1}^n b_{11}^{*l} \left[\varphi_1^*(k - m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(k - m_l) \right] \\ \quad \text{if } k \in \mathbb{Z}_0^{m_1}, \\ \\ y_1(0) = \varphi_1^*(0), \\ \\ y_2(k+1) = \lambda y_2(k) - \sum_{l=1}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(k - m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(k - m_l) \right] \\ \quad \text{if } k \in \mathbb{Z}_0^{m_1}, \\ \\ y_2(0) = \varphi_2^*(0). \end{array} \right.$$

With the aid of formula (1.20), we get

$$y_1(k) = \lambda^k \varphi_1^*(0) + \sum_{r=0}^{k-1} \lambda^{k-1-r} \left(\sum_{l=1}^n b_{11}^{*l} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_l) \right] \right),$$

$$k \in \mathbb{Z}_1^{m_1+1}, \quad (2.49)$$

$$y_2(k) = \lambda^k \varphi_2^*(0) - \sum_{r=0}^{k-1} \lambda^{k-1-r} \left(\sum_{l=1}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_l) \right] \right),$$

$$k \in \mathbb{Z}_1^{m_1+1}. \quad (2.50)$$

Now we solve system (2.47), (2.48) for $k \in \mathbb{Z}_{m_1+1}^{m_2}$, i.e., we consider the problem (with initial data deduced from (2.49), (2.50))

$$\left\{ \begin{array}{l} y_1(k+1) = \lambda y_1(k) + b_{11}^{*1} \lambda^{k-m_1} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \\ \quad + \sum_{l=2}^n b_{11}^{*l} \left[\varphi_1^*(k - m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(k - m_l) \right] \\ \quad \text{if } k \in \mathbb{Z}_{m_1+1}^{m_2}, \\ y_1(m_1+1) = \lambda^{m_1+1} \varphi_1^*(0) \\ \quad + \sum_{r=0}^{m_1} \lambda^{m_1-r} \left(\sum_{l=1}^n b_{11}^{*l} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_l) \right] \right), \\ \\ y_2(k+1) = \lambda y_2(k) - \frac{(b_{11}^{*1})^2}{b_{12}^{*1}} \lambda^{k-m_1} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \\ \quad - \sum_{l=2}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(k - m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(k - m_l) \right] \\ \quad \text{if } k \in \mathbb{Z}_{m_1+1}^{m_2}, \\ y_2(m_1+1) = \lambda^{m_1+1} \varphi_2^*(0) \\ \quad - \sum_{r=0}^{m_1} \lambda^{m_1-r} \left(\sum_{l=1}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_l) \right] \right). \end{array} \right.$$

Formula (1.20) yields (for $k \in \mathbb{Z}_{m_1+2}^{m_2+1}$)

$$\begin{aligned} y_1(k) &= \lambda^{k-(m_1+1)} y_1(m_1+1) + \sum_{r=m_1+1}^{k-1} \lambda^{k-1-r} \left(b_{11}^{*1} \lambda^{k-m_1} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right. \\ &\quad \left. + \sum_{l=2}^n b_{11}^{*l} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_l) \right] \right) \\ &= \lambda^{k-m_1-1} \left[\lambda^{m_1+1} \varphi_1^*(0) + \sum_{r=0}^{m_1} \lambda^{m_1-r} \left(\sum_{l=1}^n b_{11}^{*l} \left[\varphi_1^*(r - m_l) \right] \right) \right] \end{aligned}$$

$$\begin{aligned}
 & + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_l) \Big] \Big] + \sum_{r=m_1+1}^{k-1} \lambda^{k-1-r} \left(b_{11}^{*1} \lambda^{k-m_1} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right. \\
 & \left. + \sum_{l=2}^n b_{11}^{*l} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_l) \right] \right) \\
 & = \lambda^k \varphi_1^*(0) + \sum_{r=0}^{m_1} \lambda^{k-1-r} \left(\sum_{l=1}^n b_{11}^{*l} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_l) \right] \right) \\
 & \quad + \sum_{r=m_1+1}^{k-1} \lambda^{k-1-r} \left(b_{11}^{*1} \lambda^{k-m_1} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right. \\
 & \quad \left. + \sum_{l=2}^n b_{11}^{*l} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_l) \right] \right) \\
 & = \lambda^k \varphi_1^*(0) + \sum_{r=0}^{k-1} \lambda^{k-1-r} \left(\sum_{l=2}^n b_{11}^{*l} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_l) \right] \right) \\
 & \quad + \sum_{r=0}^{m_1} \lambda^{k-1-r} b_{11}^{*1} \left[\varphi_1^*(r - m_1) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_1) \right] \\
 & \quad + (k - 1 - m_1) b_{11}^{*1} \lambda^{k-1-m_1} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \\
 & = \lambda^k \varphi_1^*(0) + \sum_{r=0}^{k-1} \lambda^{k-1-r} \left(\sum_{l=2}^n b_{11}^{*l} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_l) \right] \right) \\
 & \quad + b_{11}^{*1} \left(\sum_{r=0}^{m_1} \lambda^{k-1-r} \left[\varphi_1^*(r - m_1) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_1) \right] \right. \\
 & \quad \left. + (k - 1 - m_1) \lambda^{k-1-m_1} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right), \tag{2.51}
 \end{aligned}$$

$$\begin{aligned}
 y_2(k) & = \lambda^{k-(m_1+1)} y_2(m_1 + 1) - \sum_{r=m_1+1}^{k-1} \lambda^{k-1-r} \left(\frac{(b_{11}^{*1})^2}{b_{12}^{*1}} \lambda^{r-m_1} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right. \\
 & \quad \left. + \sum_{l=2}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_l) \right] \right) \\
 & = \lambda^{k-m_1-1} \left[\lambda^{m_1+1} \varphi_2^*(0) - \sum_{r=0}^{m_1} \lambda^{m_1-r} \left(\sum_{l=1}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(r - m_l) \right. \right. \right. \\
 & \quad \left. \left. + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_l) \right] \right) \Big]
 \end{aligned}$$

$$\begin{aligned}
 & - \sum_{r=m_1+1}^{k-1} \lambda^{k-1-r} \left(\frac{(b_{11}^{*1})^2}{b_{12}^{*1}} \lambda^{r-m_1} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right. \\
 & \left. + \sum_{l=2}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(r-m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r-m_l) \right] \right) \\
 = & \lambda^k \varphi_2^*(0) - \sum_{r=0}^{m_1} \lambda^{k-1-r} \left(\sum_{l=1}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(r-m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r-m_l) \right] \right) \\
 & - \sum_{r=m_1+1}^{k-1} \lambda^{k-1-r} \left(\frac{(b_{11}^{*1})^2}{b_{12}^{*1}} \lambda^{r-m_1} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right. \\
 & \left. + \sum_{l=2}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(r-m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r-m_l) \right] \right) \\
 = & \lambda^k \varphi_2^*(0) - \sum_{r=0}^{k-1} \lambda^{k-1-r} \left(\sum_{l=2}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(r-m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r-m_l) \right] \right) \\
 & - \sum_{r=0}^{m_1} \lambda^{k-1-r} \frac{(b_{11}^{*1})^2}{b_{12}^{*1}} \left[\varphi_1^*(r-m_1) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r-m_1) \right] \\
 & - (k-1-m_1) \lambda^{k-1-m_1} \frac{(b_{11}^{*1})^2}{b_{12}^{*1}} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \\
 = & \lambda^k \varphi_2^*(0) - \sum_{r=0}^{k-1} \lambda^{k-1-r} \left(\sum_{l=2}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(r-m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} b_{12}^{*l} \varphi_2^*(r-m_l) \right] \right) \\
 & - \frac{(b_{11}^{*1})^2}{b_{12}^{*1}} \left(\sum_{r=0}^{m_1} \lambda^{k-1-r} \left[\varphi_1^*(r-m_1) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r-m_1) \right] \right. \\
 & \left. + (k-1-m_1) \lambda^{k-1-m_1} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right). \tag{2.52}
 \end{aligned}$$

Now we solve equations (2.47), (2.48) for $k \in \mathbb{Z}_{m_2+1}^{m_3}$, i.e., we consider the problem (with initial data deduced from (2.51), (2.52))

$$\left\{ \begin{array}{l} y_1(k+1) = \lambda y_1(k) + \sum_{l=1}^2 b_{11}^{*l} \lambda^{k-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \\ \quad + \sum_{l=3}^n b_{11}^{*l} \left[\varphi_1^*(k-m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(k-m_l) \right] \quad \text{if } k \in \mathbb{Z}_{m_2+1}^{m_3}, \\ y_1(m_2+1) = \lambda^{m_2+1} \varphi_1^*(0) \\ \quad + \sum_{r=0}^{m_2} \lambda^{m_2-r} \left(\sum_{l=2}^n b_{11}^{*l} \left[\varphi_1^*(r-m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r-m_l) \right] \right) \\ \quad + b_{11}^{*1} \left(\sum_{r=0}^{m_1} \lambda^{m_2-r} \left[\varphi_1^*(r-m_1) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r-m_1) \right] \right) \\ \quad + (m_2 - m_1) \lambda^{m_2-m_1} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \end{array} \right.$$

$$\left\{ \begin{array}{l} y_2(k+1) = \lambda y_2(k) - \sum_{l=1}^2 \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \\ \quad - \sum_{l=3}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(k-m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(k-m_l) \right] \quad \text{if } k \in \mathbb{Z}_{m_2+1}^{m_3}, \\ y_2(m_2+1) = \lambda^{m_2+1} \varphi_2^*(0) \\ \quad - \sum_{r=0}^{m_2} \lambda^{m_2-r} \left(\sum_{l=2}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(r-m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} b_{12}^{*l} \varphi_2^*(r-m_l) \right] \right) \\ \quad - \frac{(b_{11}^{*1})^2}{b_{12}^{*1}} \left(\sum_{r=0}^{m_1} \lambda^{m_2-r} \left[\varphi_1^*(r-m_1) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r-m_1) \right] \right) \\ \quad + (m_2 - m_1) \lambda^{m_2-m_1} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \end{array} \right.$$

Applying formula (1.20) yields (for $k \in \mathbb{Z}_{m_2+2}^{m_3+1}$)

$$\begin{aligned} y_1(k) &= \lambda^{k-(m_2+1)} y_1(m_2+1) + \sum_{r=m_2+1}^{k-1} \lambda^{k-1-r} \left(\sum_{l=1}^2 b_{11}^{*l} \lambda^{r-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right. \\ &\quad \left. + \sum_{l=3}^n b_{11}^{*l} \left[\varphi_1^*(r-m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r-m_l) \right] \right) \\ &= \lambda^{k-m_2-1} \left[\lambda^{m_2+1} \varphi_1^*(0) + \sum_{r=0}^{m_2} \lambda^{m_2-r} \left(\sum_{l=2}^n b_{11}^{*l} \left[\varphi_1^*(r-m_l) \right. \right. \right. \\ &\quad \left. \left. + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r-m_l) \right] \right) \end{aligned}$$

$$\begin{aligned}
 & + b_{11}^{*1} \left(\sum_{r=0}^{m_1} \lambda^{m_2-r} \left[\varphi_1^*(r - m_1) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_1) \right] \right. \\
 & \left. + (m_2 - m_1) \lambda^{m_2-m_1} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right) \\
 & + \sum_{r=m_2+1}^{k-1} \lambda^{k-1-r} \left(\sum_{l=1}^2 b_{11}^{*l} \lambda^{r-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right. \\
 & \left. + \sum_{l=3}^n b_{11}^{*l} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_l) \right] \right) \\
 & = \lambda^k \varphi_1^*(0) + \sum_{r=0}^{m_2} \lambda^{k-1-r} \left(\sum_{l=2}^n b_{11}^{*l} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_l) \right] \right) \\
 & + b_{11}^{*1} \left(\sum_{r=0}^{m_1} \lambda^{k-1-r} \left[\varphi_1^*(r - m_1) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_1) \right] \right. \\
 & \left. + (m_2 - m_1) \lambda^{k-1-m_1} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right) \\
 & + \sum_{r=m_2+1}^{k-1} \lambda^{k-1-r} \left(\sum_{l=1}^2 b_{11}^{*l} \lambda^{r-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right. \\
 & \left. + \sum_{l=3}^n b_{11}^{*l} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_l) \right] \right) \\
 & = \lambda^k \varphi_1^*(0) + \sum_{r=0}^{k-1} \lambda^{k-1-r} \left(\sum_{l=3}^n b_{11}^{*l} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_l) \right] \right) \\
 & + \sum_{r=0}^{m_2} \lambda^{k-1-r} b_{11}^{*2} \left[\varphi_1^*(r - m_2) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_2) \right] \\
 & + b_{11}^{*1} \left(\sum_{r=0}^{m_1} \lambda^{k-1-r} \left[\varphi_1^*(r - m_1) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_1) \right] \right. \\
 & \left. + (m_2 - m_1) \lambda^{k-1-m_1} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right) \\
 & + (k - 1 - m_2) \left(b_{11}^{*1} \lambda^{k-1-m_1} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right. \\
 & \left. + b_{11}^{*2} \lambda^{k-1-m_2} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right)
 \end{aligned}$$

$$\begin{aligned}
 &= \lambda^k \varphi_1^*(0) + \sum_{r=0}^{k-1} \lambda^{k-1-r} \left(\sum_{l=3}^n b_{11}^{*l} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_l) \right] \right) \\
 &\quad + b_{11}^{*1} \left(\sum_{r=0}^{m_1} \lambda^{k-1-r} \left[\varphi_1^*(r - m_1) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_1) \right] \right) \\
 &\quad + (k - 1 - m_1) \lambda^{k-1-m_1} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \\
 &\quad + b_{11}^{*2} \left(\sum_{r=0}^{m_2} \lambda^{k-1-r} \left[\varphi_1^*(r - m_2) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_2) \right] \right) \\
 &\quad + (k - 1 - m_2) \lambda^{k-1-m_2} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right], \tag{2.53}
 \end{aligned}$$

$$\begin{aligned}
 y_2(k) &= \lambda^{k-(m_2+1)} y_2(m_2 + 1) - \sum_{r=m_2+1}^{k-1} \lambda^{k-1-r} \left(\sum_{l=1}^2 \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \lambda^{r-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right) \\
 &\quad + \sum_{l=3}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_l) \right] \\
 &= \lambda^{k-m_2-1} \left[\lambda^{m_2+1} \varphi_2^*(0) - \sum_{r=0}^{m_2} \lambda^{m_2-r} \left(\sum_{l=2}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(r - m_l) \right. \right. \right. \\
 &\quad \left. \left. \left. + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_l) \right] \right) \right. \\
 &\quad \left. - \frac{(b_{11}^{*1})^2}{b_{12}^{*1}} \left(\sum_{r=0}^{m_1} \lambda^{m_2-r} \left[\varphi_1^*(r - m_1) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_1) \right] \right) \right. \\
 &\quad \left. + (m_2 - m_1) \lambda^{m_2-m_1} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right] \\
 &\quad - \sum_{r=m_2+1}^{k-1} \lambda^{k-1-r} \left(\sum_{l=1}^2 \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \lambda^{r-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right) \\
 &\quad + \sum_{l=3}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_l) \right] \\
 &= \lambda^k \varphi_2^*(0) - \sum_{r=0}^{m_2} \lambda^{k-1-r} \left(\sum_{l=2}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_l) \right] \right) \\
 &\quad - \frac{(b_{11}^{*1})^2}{b_{12}^{*1}} \left(\sum_{r=0}^{m_1} \lambda^{k-1-r} \left[\varphi_1^*(r - m_1) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_1) \right] \right)
 \end{aligned}$$

$$\begin{aligned}
 & + (m_2 - m_1) \lambda^{k-1-m_1} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \\
 & - \sum_{r=m_2+1}^{k-1} \lambda^{k-1-r} \left(\sum_{l=1}^2 \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \lambda^{r-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right. \\
 & \left. + \sum_{l=3}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_l) \right] \right) \\
 = & \lambda^k \varphi_2^*(0) - \sum_{r=0}^{k-1} \lambda^{k-1-r} \left(\sum_{l=3}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_l) \right] \right) \\
 & - \sum_{r=0}^{m_2} \lambda^{k-1-r} \frac{(b_{11}^{*2})^2}{b_{12}^{*2}} \left[\varphi_1^*(r - m_2) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_2) \right] \\
 & - \frac{(b_{11}^{*1})^2}{b_{12}^{*1}} \left(\sum_{r=0}^{m_1} \lambda^{k-1-r} \left[\varphi_1^*(r - m_1) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_1) \right] \right) \\
 & + (m_2 - m_1) \lambda^{k-1-m_1} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \\
 & - (k - 1 - m_2) \left(\frac{(b_{11}^{*1})^2}{b_{12}^{*1}} \lambda^{k-1-m_1} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right. \\
 & \left. + \frac{(b_{11}^{*2})^2}{b_{12}^{*2}} \lambda^{k-1-m_2} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right) \\
 = & \lambda^k \varphi_2^*(0) - \sum_{r=0}^{k-1} \lambda^{k-1-r} \left(\sum_{l=3}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_l) \right] \right) \\
 & - \frac{(b_{11}^{*1})^2}{b_{12}^{*1}} \left(\sum_{r=0}^{m_1} \lambda^{k-1-r} \left[\varphi_1^*(r - m_1) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_1) \right] \right) \\
 & + (k - 1 - m_1) \lambda^{k-1-m_1} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \\
 & - \frac{(b_{11}^{*2})^2}{b_{12}^{*2}} \left(\sum_{r=0}^{m_2} \lambda^{k-1-r} \left[\varphi_1^*(r - m_2) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_2) \right] \right) \\
 & + (k - 1 - m_2) \lambda^{k-1-m_2} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right]. \tag{2.54}
 \end{aligned}$$

From (2.51)–(2.54) we deduce that expected form of the solution of the initial problem for $k \in \mathbb{Z}_{m_{s-1}+1}^{m_s}$ with initial data derived from the solution of previous equation for $k \in \mathbb{Z}_{m_{s-2}+1}^{m_{s-1}}$ is

$$\begin{aligned}
 y_1(k) = & \lambda^k \varphi_1^*(0) + \sum_{r=0}^{k-1} \lambda^{k-1-r} \left(\sum_{l=s}^n b_{11}^{*l} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r - m_l) \right] \right) \\
 & + \sum_{l=1}^{s-1} \left[b_{11}^{*l} \left(\sum_{r=0}^{m_l} \lambda^{k-1-r} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r - m_l) \right] \right) \right. \\
 & \left. + (k - 1 - m_l) \lambda^{k-1-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \right) \Big] \quad (2.55)
 \end{aligned}$$

$$\text{if } k \in \mathbb{Z}_{m_{s-1}+2}^{m_s+1},$$

$$\begin{aligned}
 y_1(k) = & \lambda^k \varphi_2^*(0) - \sum_{r=0}^{k-1} \lambda^{k-1-r} \left(\sum_{l=s}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r - m_l) \right] \right) \\
 & - \sum_{l=1}^{s-1} \left[\frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left(\sum_{r=0}^{m_l} \lambda^{k-1-r} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r - m_l) \right] \right) \right. \\
 & \left. + (k - 1 - m_l) \lambda^{k-1-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \right) \Big] \quad (2.56)
 \end{aligned}$$

$$\text{if } k \in \mathbb{Z}_{m_{s-1}+2}^{m_s+1}.$$

We solve equations (2.47), (2.48) for $k \in \mathbb{Z}_{m_s+1}^{m_s+1}$ with initial data deduced from (2.55) and (2.56), i.e., we consider the problem

$$\left\{ \begin{array}{l}
 y_1(k+1) = \lambda y_1(k) + \sum_{l=1}^s b_{11}^{*l} \lambda^{k-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \\
 \quad + \sum_{l=s+1}^n b_{11}^{*l} \left[\varphi_1^*(k - m_l) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(k - m_l) \right] \\
 \quad \text{if } k \in \mathbb{Z}_{m_s+1}^{m_s+1}, \\
 y_1(m_s + 1) = \lambda^{m_s+1} \varphi_1^*(0) \\
 \quad + \sum_{r=0}^{m_s} \lambda^{m_s-r} \left(\sum_{l=s}^n b_{11}^{*l} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r - m_l) \right] \right) \\
 \quad + \sum_{l=1}^{s-1} b_{11}^{*l} \left(\sum_{r=0}^{m_l} \lambda^{m_s-r} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r - m_l) \right] \right) \\
 \quad + (m_s - m_l) \lambda^{m_s-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \Big],
 \end{array} \right.$$

$$\left\{ \begin{array}{l} y_2(k+1) = \lambda y_2(k) - \sum_{l=1}^s \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \\ \quad - \sum_{l=s+1}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(k-m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(k-m_l) \right] \\ \quad \text{if } k \in \mathbb{Z}_{m_s+1}^{m_s+1}, \\ y_2(m_s+1) = \lambda^{m_s+1} \varphi_2^*(0) \\ \quad - \sum_{r=0}^{m_s} \lambda^{m_s-r} \left(\sum_{l=s}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(r-m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r-m_l) \right] \right) \\ \quad - \sum_{l=1}^{s-1} \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left(\sum_{r=0}^{m_l} \lambda^{m_s-r} \left[\varphi_1^*(r-m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r-m_l) \right] \right) \\ \quad + (m_s - m_l) \lambda^{m_s-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \end{array} \right\}.$$

Applying formula (1.20) yields (for $k \in \mathbb{Z}_{m_s+2}^{m_s+1+1}$)

$$\begin{aligned} y_1(k) &= \lambda^{k-(m_s+1)} y_1(m_s+1) + \sum_{r=m_s+1}^{k-1} \lambda^{k-1-r} \left(\sum_{l=1}^s b_{11}^{*l} \lambda^{r-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right. \\ &\quad \left. + \sum_{l=s+1}^n b_{11}^{*l} \left[\varphi_1^*(r-m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r-m_l) \right] \right) \\ &= \lambda^{k-m_s-1} \left[\lambda^{m_s+1} \varphi_1^*(0) + \sum_{r=0}^{m_s} \lambda^{m_s-r} \left(\sum_{l=s}^n b_{11}^{*l} \left[\varphi_1^*(r-m_l) \right. \right. \right. \\ &\quad \left. \left. + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r-m_l) \right] \right) \\ &\quad \left. + \sum_{l=1}^{s-1} b_{11}^{*l} \left(\sum_{r=0}^{m_l} \lambda^{m_s-r} \left[\varphi_1^*(r-m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r-m_l) \right] \right) \right. \\ &\quad \left. + (m_s - m_l) \lambda^{m_s-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right) \\ &\quad + \sum_{r=m_s+1}^{k-1} \lambda^{k-1-r} \left(\sum_{l=1}^s b_{11}^{*l} \lambda^{r-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right. \\ &\quad \left. + \sum_{l=s+1}^n b_{11}^{*l} \left[\varphi_1^*(r-m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r-m_l) \right] \right) \end{aligned}$$

$$\begin{aligned}
 &= \lambda^k \varphi_1^*(0) + \sum_{r=0}^{m_s} \lambda^{k-1-r} \left(\sum_{l=s}^n b_{11}^{*l} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r - m_l) \right] \right) \\
 &\quad + \sum_{l=1}^{s-1} b_{11}^{*l} \left(\sum_{r=0}^{m_l} \lambda^{k-1-r} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r - m_l) \right] \right) \\
 &\quad + (m_s - m_l) \lambda^{k-1-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \\
 &\quad + \sum_{r=m_s+1}^{k-1} \lambda^{k-1-r} \left(\sum_{l=1}^s b_{11}^{*l} \lambda^{r-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \right) \\
 &\quad + \sum_{l=s+1}^n b_{11}^{*l} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r - m_l) \right] \\
 &= \lambda^k \varphi_1^*(0) + \sum_{r=0}^{k-1} \lambda^{k-1-r} \left(\sum_{l=s+1}^n b_{11}^{*l} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r - m_l) \right] \right) \\
 &\quad + b_{11}^{*s} \sum_{r=0}^{m_s} \lambda^{k-1-r} \left[\varphi_1^*(r - m_s) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r - m_s) \right] \\
 &\quad + b_{11}^{*1} \left(\sum_{r=0}^{m_1} \lambda^{k-1-r} \left[\varphi_1^*(r - m_1) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r - m_1) \right] \right) \\
 &\quad + (m_s - m_1) \lambda^{k-1-m_1} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \\
 &\quad + b_{11}^{*2} \left(\sum_{r=0}^{m_2} \lambda^{k-1-r} \left[\varphi_1^*(r - m_2) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r - m_2) \right] \right) \\
 &\quad + (m_s - m_2) \lambda^{k-1-m_2} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \\
 &\quad + \dots \\
 &\quad + b_{11}^{*s-1} \left(\sum_{r=0}^{m_{s-1}} \lambda^{k-1-r} \left[\varphi_1^*(r - m_{s-1}) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r - m_{s-1}) \right] \right) \\
 &\quad + (m_s - m_{s-1}) \lambda^{k-1-m_{s-1}} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \\
 &\quad + (k - 1 - m_s) \left(\lambda^{k-1-m_1} b_{11}^{*1} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \right) \\
 &\quad + \lambda^{k-1-m_2} b_{11}^{*2} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right]
 \end{aligned}$$

$$\begin{aligned}
 & + \dots \\
 & + \lambda^{k-1-m_{s-1}} b_{11}^{*s-1} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \\
 & + \lambda^{k-1-m_s} b_{11}^{*s} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \\
 & = \lambda^k \varphi_1^*(0) + \sum_{r=0}^{k-1} \lambda^{k-1-r} \left(\sum_{l=s+1}^n b_{11}^{*l} \left[\varphi_1^*(r-m_l) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r-m_l) \right] \right) \\
 & + b_{11}^{*1} \left(\sum_{r=0}^{m_1} \lambda^{k-1-r} \left[\varphi_1^*(r-m_1) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r-m_1) \right] \right) \\
 & + (k-1-m_1) \lambda^{k-1-m_1} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \\
 & + b_{11}^{*2} \left(\sum_{r=0}^{m_2} \lambda^{k-1-r} \left[\varphi_1^*(r-m_2) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r-m_2) \right] \right) \\
 & + (k-1-m_2) \lambda^{k-1-m_2} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \\
 & + \dots \\
 & + b_{11}^{*s-1} \left(\sum_{r=0}^{m_{s-1}} \lambda^{k-1-r} \left[\varphi_1^*(r-m_{s-1}) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r-m_{s-1}) \right] \right) \\
 & + (k-1-m_{s-1}) \lambda^{k-1-m_{s-1}} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \\
 & + b_{11}^{*s} \left(\sum_{r=0}^{m_s} \lambda^{k-1-r} \left[\varphi_1^*(r-m_s) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r-m_s) \right] \right) \\
 & + (k-1-m_s) \lambda^{k-1-m_s} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \\
 & = \lambda^k \varphi_1^*(0) + \sum_{r=0}^{k-1} \lambda^{k-1-r} \left(\sum_{l=s+1}^n b_{11}^{*l} \left[\varphi_1^*(r-m_l) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r-m_l) \right] \right) \\
 & + \sum_{l=1}^s b_{11}^{*l} \left(\sum_{r=0}^{m_l} \lambda^{k-1-r} \left[\varphi_1^*(r-m_l) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r-m_l) \right] \right) \\
 & + (k-1-m_l) \lambda^{k-1-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right], \tag{2.57}
 \end{aligned}$$

$$\begin{aligned}
 y_2(k) &= \lambda^{k-(m_s+1)} y_2(m_s+1) - \sum_{r=m_s+1}^{k-1} \lambda^{k-1-r} \left(\sum_{l=1}^s \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \lambda^{r-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right. \\
 &\quad \left. + \sum_{l=s+1}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(r-m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r-m_l) \right] \right) \\
 &= \lambda^{k-m_s-1} \left[\lambda^{m_s+1} \varphi_2^*(0) - \sum_{r=0}^{m_s} \lambda^{m_s-r} \left(\sum_{l=s}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(r-m_l) \right. \right. \right. \\
 &\quad \left. \left. \left. + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r-m_l) \right] \right) \right. \\
 &\quad \left. - \sum_{l=1}^{s-1} \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left(\sum_{r=0}^{m_l} \lambda^{m_s-r} \left[\varphi_1^*(r-m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r-m_l) \right] \right) \right. \\
 &\quad \left. + (m_s - m_l) \lambda^{m_s-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right) \\
 &\quad - \sum_{r=m_s+1}^{k-1} \lambda^{k-1-r} \left(\sum_{l=1}^s \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \lambda^{r-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right. \\
 &\quad \left. + \sum_{l=s+1}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(r-m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r-m_l) \right] \right) \\
 &= \lambda^k \varphi_2^*(0) - \sum_{r=0}^{m_s} \lambda^{k-1-r} \left(\sum_{l=s}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(r-m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r-m_l) \right] \right) \\
 &\quad - \sum_{l=1}^{s-1} \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left(\sum_{r=0}^{m_l} \lambda^{k-1-r} \left[\varphi_1^*(r-m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r-m_l) \right] \right. \\
 &\quad \left. + (m_s - m_l) \lambda^{k-1-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right) \\
 &\quad - \sum_{r=m_s+1}^{k-1} \lambda^{k-1-r} \left(\sum_{l=1}^s \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \lambda^{r-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right. \\
 &\quad \left. + \sum_{l=s+1}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(r-m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r-m_l) \right] \right) \\
 &= \lambda^k \varphi_2^*(0) - \sum_{r=0}^{k-1} \lambda^{k-1-r} \left(\sum_{l=s+1}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(r-m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r-m_l) \right] \right) \\
 &\quad - \frac{(b_{11}^{*s})^2}{b_{12}^{*s}} \sum_{r=0}^{m_s} \lambda^{k-1-r} \left[\varphi_1^*(r-m_s) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r-m_s) \right]
 \end{aligned}$$

$$\begin{aligned}
 & - \frac{(b_{11}^{*1})^2}{b_{12}^{*1}} \left(\sum_{r=0}^{m_1} \lambda^{k-1-r} \left[\varphi_1^*(r - m_1) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_1) \right] \right. \\
 & \left. + (m_s - m_1) \lambda^{k-1-m_1} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right) \\
 & - \frac{(b_{11}^{*2})^2}{b_{12}^{*2}} \left(\sum_{r=0}^{m_2} \lambda^{k-1-r} \left[\varphi_1^*(r - m_2) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_2) \right] \right. \\
 & \left. + (m_s - m_2) \lambda^{k-1-m_2} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right) \\
 & + \dots \\
 & - \frac{(b_{11}^{*s-1})^2}{b_{12}^{*s-1}} \left(\sum_{r=0}^{m_{s-1}} \lambda^{k-1-r} \left[\varphi_1^*(r - m_{s-1}) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_{s-1}) \right] \right. \\
 & \left. + (m_s - m_{s-1}) \lambda^{k-1-m_{s-1}} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right) \\
 & - (k - 1 - m_s) \left(\lambda^{k-1-m_1} \frac{(b_{11}^{*1})^2}{b_{12}^{*1}} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right. \\
 & \left. + \lambda^{k-1-m_2} \frac{(b_{11}^{*2})^2}{b_{12}^{*2}} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] + \dots \right. \\
 & \left. + \lambda^{k-1-m_{s-1}} \frac{(b_{11}^{*s-1})^2}{b_{12}^{*s-1}} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right. \\
 & \left. + \lambda^{k-1-m_s} \frac{(b_{11}^{*s})^2}{b_{12}^{*s}} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right) \\
 & = \lambda^k \varphi_2^*(0) - \sum_{r=0}^{k-1} \lambda^{k-1-r} \left(\sum_{l=s+1}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_l) \right] \right) \\
 & - \frac{(b_{11}^{*1})^2}{b_{12}^{*1}} \left(\sum_{r=0}^{m_1} \lambda^{k-1-r} \left[\varphi_1^*(r - m_1) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_1) \right] \right. \\
 & \left. + (k - 1 - m_1) \lambda^{k-1-m_1} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right) \\
 & - \frac{(b_{11}^{*2})^2}{b_{12}^{*2}} \left(\sum_{r=0}^{m_2} \lambda^{k-1-r} \left[\varphi_1^*(r - m_2) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_2) \right] \right. \\
 & \left. + (k - 1 - m_2) \lambda^{k-1-m_2} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right)
 \end{aligned}$$

$$\begin{aligned}
 & + \dots \\
 & - \frac{(b_{11}^{*s-1})^2}{b_{12}^{*s-1}} \left(\sum_{r=0}^{m_{s-1}} \lambda^{k-1-r} \left[\varphi_1^*(r - m_{s-1}) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r - m_{s-1}) \right] \right. \\
 & \left. + (k - 1 - m_{s-1}) \lambda^{k-1-m_{s-1}} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \right) \\
 & - \frac{(b_{11}^{*s})^2}{b_{12}^{*s}} \left(\sum_{r=0}^{m_s} \lambda^{k-1-r} \left[\varphi_1^*(r - m_s) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r - m_s) \right] \right. \\
 & \left. + (k - 1 - m_s) \lambda^{k-1-m_s} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \right) \\
 & = \lambda^k \varphi_2^*(0) - \sum_{r=0}^{k-1} \lambda^{k-1-r} \left(\sum_{l=s+1}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r - m_l) \right] \right) \\
 & - \sum_{l=1}^s \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left(\sum_{r=0}^{m_l} \lambda^{k-1-r} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r - m_l) \right] \right. \\
 & \left. + (k - 1 - m_l) \lambda^{k-1-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \right). \tag{2.58}
 \end{aligned}$$

In the end, we solve equations (2.47), (2.48) for $k \in \mathbb{Z}_{m_n+1}^\infty$ with initial data deduced from (2.57) and (2.58), i.e., we consider the problem

$$\left\{ \begin{array}{l}
 y_1(k+1) = \lambda y_1(k) + \sum_{l=1}^n b_{11}^{*l} \lambda^{k-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \quad \text{if } k \in \mathbb{Z}_{m_n+1}^\infty, \\
 y_1(m_n+1) = \lambda^{m_n+1} \varphi_1^*(0) + \sum_{r=0}^{m_n} \lambda^{m_n-r} b_{11}^{*n} \left[\varphi_1^*(r - m_n) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r - m_n) \right] \\
 \quad + \sum_{l=1}^{n-1} b_{11}^{*l} \left(\sum_{r=0}^{m_l} \lambda^{m_n-r} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r - m_l) \right] \right. \\
 \quad \left. + (m_n - m_l) \lambda^{m_n-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \right),
 \end{array} \right.$$

$$\left\{ \begin{array}{l} y_2(k+1) = \lambda y_2(k) - \sum_{l=1}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \lambda^{k-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \text{ if } k \in \mathbb{Z}_{m_n+1}^\infty, \\ y_2(m_n+1) = \lambda^{m_n+1} \varphi_2^*(0) \\ \quad - \sum_{r=0}^{m_n} \lambda^{m_n-r} \frac{(b_{11}^{*n})^2}{b_{12}^{*n}} \left[\varphi_1^*(r-m_n) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r-m_n) \right] \\ \quad - \sum_{l=1}^{n-1} \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left(\sum_{r=0}^{m_l} \lambda^{m_n-r} \left[\varphi_1^*(r-m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r-m_l) \right] \right. \\ \quad \left. + (m_n - m_l) \lambda^{m_n-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right). \end{array} \right.$$

Applying formula (1.20) yields (for $k \in \mathbb{Z}_{m_n+2}^\infty$)

$$\begin{aligned} y_1(k) &= \lambda^{k-(m_n+1)} y_1(m_n+1) + \sum_{r=m_n+1}^{k-1} \lambda^{k-1-r} \left(\sum_{l=1}^n b_{11}^{*l} \lambda^{r-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right) \\ &= \lambda^{k-m_n-1} \left[\lambda^{m_n+1} \varphi_1^*(0) + \sum_{r=0}^{m_n} \lambda^{m_n-r} b_{11}^{*n} \left[\varphi_1^*(r-m_n) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r-m_n) \right] \right. \\ &\quad + \sum_{l=1}^{n-1} b_{11}^{*l} \left(\sum_{r=0}^{m_l} \lambda^{m_n-r} \left[\varphi_1^*(r-m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r-m_l) \right] \right. \\ &\quad \left. \left. + (m_n - m_l) \lambda^{m_n-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right) \right] \\ &\quad + \sum_{r=m_n+1}^{k-1} \lambda^{k-1-r} \left(\sum_{l=1}^n b_{11}^{*l} \lambda^{r-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right) \\ &= \lambda^k \varphi_1^*(0) + \sum_{r=0}^{m_n} \lambda^{k-1-r} b_{11}^{*n} \left[\varphi_1^*(r-m_n) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r-m_n) \right] \\ &\quad + \sum_{l=1}^{n-1} b_{11}^{*l} \left(\sum_{r=0}^{m_l} \lambda^{k-1-r} \left[\varphi_1^*(r-m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r-m_l) \right] \right. \\ &\quad \left. + (m_n - m_l) \lambda^{k-1-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right) \\ &\quad + \sum_{r=m_n+1}^{k-1} \lambda^{k-1-r} \left(\sum_{l=1}^n b_{11}^{*l} \lambda^{r-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \right) \end{aligned}$$

$$\begin{aligned}
 &= \lambda^k \varphi_1^*(0) + \sum_{r=0}^{m_n} \lambda^{k-1-r} b_{11}^{*n} \left[\varphi_1^*(r - m_n) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r - m_n) \right] \\
 &\quad + b_{11}^{*1} \left(\sum_{r=0}^{m_1} \lambda^{k-1-r} \left[\varphi_1^*(r - m_1) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r - m_1) \right] \right. \\
 &\quad \left. + (m_n - m_1) \lambda^{k-1-m_1} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \right) \\
 &\quad + b_{11}^{*2} \left(\sum_{r=0}^{m_2} \lambda^{k-1-r} \left[\varphi_1^*(r - m_2) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r - m_2) \right] \right. \\
 &\quad \left. + (m_n - m_2) \lambda^{k-1-m_2} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \right) \\
 &\quad + \dots \\
 &\quad + b_{11}^{*n-1} \left(\sum_{r=0}^{m_{n-1}} \lambda^{k-1-r} \left[\varphi_1^*(r - m_{n-1}) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r - m_{n-1}) \right] \right. \\
 &\quad \left. + (m_n - m_{n-1}) \lambda^{k-1-m_{n-1}} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \right) \\
 &\quad + (k - 1 - m_n) \left(\lambda^{k-1-m_1} b_{11}^{*1} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \right. \\
 &\quad \left. + \lambda^{k-1-m_2} b_{11}^{*2} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \right) \\
 &\quad + \dots \\
 &\quad + \lambda^{k-1-m_{n-1}} b_{11}^{*n-1} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \\
 &\quad + \lambda^{k-1-m_n} b_{11}^{*n} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \\
 &= \lambda^k \varphi_1^*(0) + b_{11}^{*1} \left(\sum_{r=0}^{m_1} \lambda^{k-1-r} \left[\varphi_1^*(r - m_1) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r - m_1) \right] \right. \\
 &\quad \left. + (k - 1 - m_1) \lambda^{k-1-m_1} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \right) \\
 &\quad + b_{11}^{*2} \left(\sum_{r=0}^{m_2} \lambda^{k-1-r} \left[\varphi_1^*(r - m_2) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r - m_2) \right] \right.
 \end{aligned}$$

$$\begin{aligned}
 & + (k-1-m_2)\lambda^{k-1-m_2} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \\
 & + \dots \\
 & + b_{11}^{*n-1} \left(\sum_{r=0}^{m_{n-1}} \lambda^{k-1-r} \left[\varphi_1^*(r-m_{n-1}) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r-m_{n-1}) \right] \right. \\
 & \left. + (k-1-m_{n-1})\lambda^{k-1-m_{n-1}} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \right) \\
 & + b_{11}^{*n} \left(\sum_{r=0}^{m_n} \lambda^{k-1-r} \left[\varphi_1^*(r-m_n) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r-m_n) \right] \right. \\
 & \left. + (k-1-m_n)\lambda^{k-1-m_n} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \right) \\
 & = \lambda^k \varphi_1^*(0) + \sum_{l=1}^n b_{11}^{*l} \left(\sum_{r=0}^{m_l} \lambda^{k-1-r} \left[\varphi_1^*(r-m_l) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r-m_l) \right] \right. \\
 & \left. + (k-1-m_l)\lambda^{k-1-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \right), \tag{2.59}
 \end{aligned}$$

$$\begin{aligned}
 y_2(k) & = \lambda^{k-(m_n+1)} y_2(m_n+1) - \sum_{r=m_n+1}^{k-1} \lambda^{k-1-r} \left(\sum_{l=1}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \lambda^{r-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \right) \\
 & = \lambda^{k-m_n-1} \left[\lambda^{m_n+1} \varphi_2^*(0) - \sum_{r=0}^{m_n} \lambda^{m_n-r} \frac{(b_{11}^{*n})^2}{b_{12}^{*n}} \left[\varphi_1^*(r-m_n) \right. \right. \\
 & \quad \left. \left. + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r-m_n) \right] \right. \\
 & \quad \left. - \sum_{l=1}^{n-1} \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left(\sum_{r=0}^{m_l} \lambda^{m_n-r} \left[\varphi_1^*(r-m_l) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r-m_l) \right] \right. \right. \\
 & \quad \left. \left. + (m_n-m_l)\lambda^{m_n-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \right) \right] \\
 & \quad - \sum_{r=m_n+1}^{k-1} \lambda^{k-1-r} \left(\sum_{l=1}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \lambda^{r-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \right) \\
 & = \lambda^k \varphi_2^*(0) - \sum_{r=0}^{m_n} \lambda^{k-1-r} \frac{(b_{11}^{*n})^2}{b_{12}^{*n}} \left[\varphi_1^*(r-m_n) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r-m_n) \right] \\
 & \quad - \sum_{l=1}^{n-1} \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left(\sum_{r=0}^{m_l} \lambda^{k-1-r} \left[\varphi_1^*(r-m_l) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r-m_l) \right] \right)
 \end{aligned}$$

$$\begin{aligned}
 & + (m_n - m_l)\lambda^{k-1-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}}\varphi_2^*(0) \right] \\
 & - \sum_{r=m_n+1}^{k-1} \lambda^{k-1-r} \left(\sum_{l=1}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \lambda^{r-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}}\varphi_2^*(0) \right] \right) \\
 = & \lambda^k \varphi_2^*(0) - \sum_{r=0}^{m_n} \lambda^{k-1-r} \frac{(b_{11}^{*n})^2}{b_{12}^{*n}} \left[\varphi_1^*(r - m_n) + \frac{b_{12}^{*1}}{b_{11}^{*1}}\varphi_2^*(r - m_n) \right] \\
 & - \frac{(b_{11}^{*1})^2}{b_{12}^{*1}} \left(\sum_{r=0}^{m_1} \lambda^{k-1-r} \left[\varphi_1^*(r - m_1) + \frac{b_{12}^{*1}}{b_{11}^{*1}}\varphi_2^*(r - m_1) \right] \right) \\
 & + (m_n - m_1)\lambda^{k-1-m_1} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}}\varphi_2^*(0) \right] \\
 & - \frac{(b_{11}^{*2})^2}{b_{12}^{*2}} \left(\sum_{r=0}^{m_2} \lambda^{k-1-r} \left[\varphi_1^*(r - m_2) + \frac{b_{12}^{*1}}{b_{11}^{*1}}\varphi_2^*(r - m_2) \right] \right) \\
 & + (m_n - m_2)\lambda^{k-1-m_2} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}}\varphi_2^*(0) \right] \\
 & + \dots \\
 & - \frac{(b_{11}^{*n-1})^2}{b_{12}^{*n-1}} \left(\sum_{r=0}^{m_{n-1}} \lambda^{k-1-r} \left[\varphi_1^*(r - m_{n-1}) + \frac{b_{12}^{*1}}{b_{11}^{*1}}\varphi_2^*(r - m_{n-1}) \right] \right) \\
 & + (m_n - m_{n-1})\lambda^{k-1-m_{n-1}} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}}\varphi_2^*(0) \right] \\
 & - (k - 1 - m_n) \left(\lambda^{k-1-m_1} \frac{(b_{11}^{*1})^2}{b_{12}^{*1}} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}}\varphi_2^*(0) \right] \right) \\
 & + \lambda^{k-1-m_2} \frac{(b_{11}^{*2})^2}{b_{12}^{*2}} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}}\varphi_2^*(0) \right] \\
 & + \dots \\
 & + \lambda^{k-1-m_{n-1}} \frac{(b_{11}^{*n-1})^2}{b_{12}^{*n-1}} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}}\varphi_2^*(0) \right] \\
 & + \lambda^{k-1-m_n} \frac{(b_{11}^{*n})^2}{b_{12}^{*n}} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}}\varphi_2^*(0) \right] \\
 = & \lambda^k \varphi_2^*(0) - \frac{(b_{11}^{*1})^2}{b_{12}^{*1}} \left(\sum_{r=0}^{m_1} \lambda^{k-1-r} \left[\varphi_1^*(r - m_1) + \frac{b_{12}^{*1}}{b_{11}^{*1}}\varphi_2^*(r - m_1) \right] \right)
 \end{aligned}$$

$$\begin{aligned}
 & + (k-1-m_1)\lambda^{k-1-m_1} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \\
 & - \frac{(b_{11}^*)^2}{b_{12}^*} \left(\sum_{r=0}^{m_2} \lambda^{k-1-r} \left[\varphi_1^*(r-m_2) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r-m_2) \right] \right) \\
 & + (k-1-m_2)\lambda^{k-1-m_2} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \\
 & + \dots \\
 & - \frac{(b_{11}^*)^2}{b_{12}^*} \left(\sum_{r=0}^{m_{n-1}} \lambda^{k-1-r} \left[\varphi_1^*(r-m_{n-1}) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r-m_{n-1}) \right] \right) \\
 & + (k-1-m_{n-1})\lambda^{k-1-m_{n-1}} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \\
 & - \frac{(b_{11}^*)^2}{b_{12}^*} \left(\sum_{r=0}^{m_n} \lambda^{k-1-r} \left[\varphi_1^*(r-m_n) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r-m_n) \right] \right) \\
 & + (k-1-m_n)\lambda^{k-1-m_n} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \\
 & = \lambda^k \varphi_2^*(0) - \sum_{l=1}^n \frac{(b_{11}^*)^2}{b_{12}^*} \left(\sum_{r=0}^{m_l} \lambda^{k-1-r} \left[\varphi_1^*(r-m_l) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r-m_l) \right] \right) \\
 & + (k-1-m_l)\lambda^{k-1-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right]. \tag{2.60}
 \end{aligned}$$

Summing up all particular cases (2.49), (2.51), (2.53), (2.57) and (2.59) we have

$$y_1(k) = \left\{ \begin{array}{l}
 \varphi_1^*(k) \quad \text{if } k \in \mathbb{Z}_{-m_n}^0, \\
 \lambda^k \varphi_1^*(0) + \sum_{r=0}^{k-1} \lambda^{k-1-r} \left(\sum_{l=1}^n b_{11}^{*l} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r - m_l) \right] \right) \\
 \quad \text{if } k \in \mathbb{Z}_{m_1+1}^{m_1+1}, \\
 \lambda^k \varphi_1^*(0) + \sum_{r=0}^{k-1} \lambda^{k-1-r} \left(\sum_{l=2}^n b_{11}^{*l} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r - m_l) \right] \right) \\
 \quad + b_{11}^{*1} \left(\sum_{r=0}^{m_1} \lambda^{k-1-r} \left[\varphi_1^*(r - m_1) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r - m_1) \right] \right) \\
 \quad + (k - 1 - m_1) \lambda^{k-1-m_1} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \\
 \quad \text{if } k \in \mathbb{Z}_{m_1+2}^{m_2+1}, \\
 \lambda^k \varphi_1^*(0) + \sum_{r=0}^{k-1} \lambda^{k-1-r} \left(\sum_{l=3}^n b_{11}^{*l} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r - m_l) \right] \right) \\
 \quad + b_{11}^{*1} \left(\sum_{r=0}^{m_1} \lambda^{k-1-r} \left[\varphi_1^*(r - m_1) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r - m_1) \right] \right) \\
 \quad + (k - 1 - m_1) \lambda^{k-1-m_1} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \\
 \quad + b_{11}^{*2} \left(\sum_{r=0}^{m_2} \lambda^{k-1-r} \left[\varphi_1^*(r - m_2) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r - m_2) \right] \right) \\
 \quad + (k - 1 - m_2) \lambda^{k-1-m_2} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \\
 \quad \text{if } k \in \mathbb{Z}_{m_2+2}^{m_3+1}, \\
 \dots \\
 \lambda^k \varphi_1^*(0) + \sum_{r=0}^{k-1} \lambda^{k-1-r} \left(\sum_{l=s+1}^n b_{11}^{*l} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r - m_l) \right] \right) \\
 \quad + \sum_{l=1}^s b_{11}^{*l} \left(\sum_{r=0}^{m_l} \lambda^{k-1-r} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r - m_l) \right] \right) \\
 \quad + (k - 1 - m_l) \lambda^{k-1-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \\
 \quad \text{if } k \in \mathbb{Z}_{m_s+2}^{m_{s+1}+1}, \\
 \dots \\
 \lambda^k \varphi_1^*(0) + \sum_{l=1}^n b_{11}^{*l} \left(\sum_{r=0}^{m_l} \lambda^{k-1-r} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(r - m_l) \right] \right) \\
 \quad + (k - 1 - m_l) \lambda^{k-1-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^*}{b_{11}^*} \varphi_2^*(0) \right] \\
 \quad \text{if } k \in \mathbb{Z}_{m_n+2}^\infty,
 \end{array} \right. \quad (2.61)$$

and from cases (2.50), (2.52), (2.54), (2.58) and (2.60) we conclude

$$y_2(k) = \left\{ \begin{array}{l} \varphi_2^*(k) \quad \text{if } k \in \mathbb{Z}_{-m_n}^0, \\ \lambda^k \varphi_2^*(0) - \sum_{r=0}^{k-1} \lambda^{k-1-r} \left(\sum_{l=1}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_l) \right] \right) \\ \quad \text{if } k \in \mathbb{Z}_1^{m_1+1}, \\ \lambda^k \varphi_2^*(0) - \sum_{r=0}^{k-1} \lambda^{k-1-r} \left(\sum_{l=2}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} b_{12}^{*l} \varphi_2^*(r - m_l) \right] \right) \\ \quad - \frac{(b_{11}^{*1})^2}{b_{12}^{*1}} \left(\sum_{r=0}^{m_1} \lambda^{k-1-r} \left[\varphi_1^*(r - m_1) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_1) \right] \right) \\ \quad + (k - 1 - m_1) \lambda^{k-1-m_1} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \\ \quad \text{if } k \in \mathbb{Z}_{m_1+2}^{m_2+1}, \\ \lambda^k \varphi_2^*(0) - \sum_{r=0}^{k-1} \lambda^{k-1-r} \left(\sum_{l=3}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_l) \right] \right) \\ \quad - \frac{(b_{11}^{*1})^2}{b_{12}^{*1}} \left(\sum_{r=0}^{m_1} \lambda^{k-1-r} \left[\varphi_1^*(r - m_1) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_1) \right] \right) \\ \quad + (k - 1 - m_1) \lambda^{k-1-m_1} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \\ \quad - \frac{(b_{11}^{*2})^2}{b_{12}^{*2}} \left(\sum_{r=0}^{m_2} \lambda^{k-1-r} \left[\varphi_1^*(r - m_2) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_2) \right] \right) \\ \quad + (k - 1 - m_2) \lambda^{k-1-m_2} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \\ \quad \text{if } k \in \mathbb{Z}_{m_2+2}^{m_3+1}, \\ \dots \\ \lambda^k \varphi_2^*(0) \sum_{r=0}^{k-1} \lambda^{k-1-r} \left(\sum_{l=s+1}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_l) \right] \right) \\ \quad - \sum_{l=1}^s \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left(\sum_{r=0}^{m_l} \lambda^{k-1-r} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_l) \right] \right) \\ \quad + (k - 1 - m_l) \lambda^{k-1-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \\ \quad \text{if } k \in \mathbb{Z}_{m_s+2}^{m_{s+1}+1}, \\ \dots \\ \lambda^k \varphi_2^*(0) - \sum_{l=1}^n \frac{(b_{11}^{*l})^2}{b_{12}^{*l}} \left(\sum_{r=0}^{m_l} \lambda^{k-1-r} \left[\varphi_1^*(r - m_l) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(r - m_l) \right] \right) \\ \quad + (k - 1 - m_l) \lambda^{k-1-m_l} \left[\varphi_1^*(0) + \frac{b_{12}^{*1}}{b_{11}^{*1}} \varphi_2^*(0) \right] \\ \quad \text{if } k \in \mathbb{Z}_{m_n+2}^\infty. \end{array} \right. \tag{2.62}$$

Formula (2.42) is now a direct consequence of (2.61), (2.62) and (2.41). \square

2.1.4 The case (2.5) of a double real root

If the matrix Λ has the form (2.5), the necessary and sufficient conditions (1.11)–(1.14), for (2.6) are reduced to (2.9), (2.10), (2.12) and

$$\begin{vmatrix} \lambda & 1 \\ b_{21}^{*l} & b_{22}^{*l} \end{vmatrix} + \begin{vmatrix} b_{11}^{*l} & b_{12}^{*l} \\ 0 & \lambda \end{vmatrix} = \lambda(b_{11}^{*l} + b_{22}^{*l}) - b_{21}^{*l} = 0. \quad (2.63)$$

Then (2.9), (2.10), and (2.63) give $b_{11}^{*l} = b_{22}^{*l} = b_{21}^{*l} = 0$.

Theorem 2.1.5. *Let (1.1) be a weakly delayed system, equation (2.1) has a double root $\lambda_{1,2} = \lambda$ and the matrix Λ has the form (2.5). Then $b_{11}^{*l} = b_{22}^{*l} = b_{21}^{*l} = 0$ and the solution of the initial problem (1.1), (1.4) is $x(k) = \mathcal{S}y(k)$, $y(k) = (y_1(k), y_2(k))^T$ and*

$$y_1(k) = \begin{cases} \varphi_1^*(k) & \text{if } k \in \mathbb{Z}_{-m_n}^0, \\ \lambda^k \varphi_1^*(0) + k\lambda^{k-1} \varphi_2^*(0) + \sum_{r=0}^{k-1} \lambda^{k-1-r} \left[\sum_{l=1}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right] \\ & \text{if } k \in \mathbb{Z}_1^{m_1+1}, \\ \dots \\ \lambda^k \varphi_1^*(0) + k\lambda^{k-1} \varphi_2^*(0) + \sum_{r=0}^{k-1} \lambda^{k-1-r} \left[\sum_{l=s+1}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right] \\ & + \sum_{l=1}^s b_{12}^{*l} \left[\sum_{r=0}^{m_l} \lambda^{k-1-r} \varphi_2^*(r - m_l) \right. \\ & \left. + (k-1-m_l) \lambda^{k-1-m_l} \varphi_2^*(0) \right] \\ & \text{if } k \in \mathbb{Z}_{m_s+2}^{m_{s+1}+1}, \quad s = 1, 2, \dots, n-1, \\ \dots \\ \lambda^k \varphi_1^*(0) + k\lambda^{k-1} \varphi_2^*(0) + \sum_{l=1}^n b_{12}^{*l} \left[\sum_{r=0}^{m_l} \lambda^{k-1-r} \varphi_2^*(r - m_l) \right. \\ & \left. + (k-1-m_l) \lambda^{k-1-m_l} \varphi_2^*(0) \right] \\ & \text{if } k \in \mathbb{Z}_{m_n+2}^\infty, \end{cases} \quad (2.64)$$

$$y_2(k) = \begin{cases} \varphi_2^*(k) & \text{if } k \in \mathbb{Z}_{-m_n}^0, \\ \lambda^k \varphi_2^*(0) & \text{if } k \in \mathbb{Z}_1^\infty. \end{cases} \quad (2.65)$$

Proof. The system (2.6) can be written as

$$y_1(k+1) = \lambda y_1(k) + y_2(k) + \sum_{l=1}^n b_{12}^{*l} y_2(k-m_l), \quad (2.66)$$

$$y_2(k+1) = \lambda y_2(k), \quad k \in \mathbb{Z}_0^\infty. \quad (2.67)$$

Solving (2.67), we get

$$y_2(k) = \begin{cases} \varphi_2^*(k) & \text{if } k \in \mathbb{Z}_{-m_n}^0, \\ \lambda^k \varphi_2^*(0) & \text{if } k \in \mathbb{Z}_1^\infty. \end{cases} \quad (2.68)$$

Then (2.66) turns into

$$y_1(k+1) = \begin{cases} \lambda y_1(k) + \lambda^k \varphi_2^*(0) + \sum_{l=1}^n b_{12}^{*l} \varphi_2^*(k-m_l) \\ \quad \text{if } k \in \mathbb{Z}_0^{m_1}, \\ \lambda y_1(k) + \lambda^k \varphi_2^*(0) + b_{12}^{*1} \lambda^{k-m_1} \varphi_2^*(0) + \sum_{l=2}^n b_{12}^{*l} \varphi_2^*(k-m_l) \\ \quad \text{if } k \in \mathbb{Z}_{m_1+1}^{m_2}, \\ \lambda y_1(k) + \lambda^k \varphi_2^*(0) + \sum_{l=1}^2 b_{12}^{*l} \lambda^{k-m_l} \varphi_2^*(0) + \sum_{l=3}^n b_{12}^{*l} \varphi_2^*(k-m_l) \\ \quad \text{if } k \in \mathbb{Z}_{m_2+1}^{m_3}, \\ \dots \\ \lambda y_1(k) + \lambda^k \varphi_2^*(0) + \sum_{l=1}^s b_{12}^{*l} \lambda^{k-m_l} \varphi_2^*(0) \\ \quad + \sum_{l=s+1}^n b_{12}^{*l} \varphi_2^*(k-m_l) \\ \quad \text{if } k \in \mathbb{Z}_{m_s+1}^{m_{s+1}}, \quad s = 3, 4, \dots, n-1, \\ \dots \\ \lambda y_1(k) + \lambda^k \varphi_2^*(0) + \sum_{l=1}^n b_{12}^{*l} \lambda^{k-m_l} \varphi_2^*(0) \\ \quad \text{if } k \in \mathbb{Z}_{m_n+1}^\infty. \end{cases} \quad (2.69)$$

The equation (2.69) can be solved in a way similar to that of equation (2.20) in the proof of Theorem 2.1.1 using (1.20).

First we solve the equation (2.69) for $k \in \mathbb{Z}_0^{m_1}$. This means that we consider the problem

$$\begin{cases} y_1(k+1) = \lambda y_1(k) + \lambda^k \varphi_2^*(0) + \sum_{l=1}^n b_{12}^{*l} \varphi_2^*(k-m_l) & \text{if } k \in \mathbb{Z}_0^{m_1}, \\ y_1(0) = \varphi_1^*(0). \end{cases}$$

With the aid of formula (1.20), we get

$$\begin{aligned} y_1(k) &= \lambda^k \varphi_1^*(0) + \sum_{r=0}^{k-1} \lambda^{k-1-r} \left[\lambda^r \varphi_2^*(0) + \sum_{l=1}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right] \\ &= \lambda^k \varphi_1^*(0) + k \lambda^{k-1} \varphi_2^*(0) + \sum_{r=0}^{k-1} \lambda^{k-1-r} \left[\sum_{l=1}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right], \quad k \in \mathbb{Z}_{m_1+1}^{m_1+1}. \end{aligned} \quad (2.70)$$

Now we solve equation (2.69) for $k \in \mathbb{Z}_{m_1+1}^{m_2}$ with initial data deduced from (2.70), i.e., we consider the problem

$$\left\{ \begin{array}{l} y_1(k+1) = \lambda y_1(k) + \lambda^k \varphi_2^*(0) + b_{12}^{*1} \lambda^{k-m_1} \varphi_2^*(0) + \sum_{l=2}^n b_{12}^{*l} \varphi_2^*(k - m_l), \\ \quad k \in \mathbb{Z}_{m_1+1}^{m_2}, \\ y_1(m_1+1) = \lambda^{m_1+1} \varphi_1^*(0) + (m_1+1) \lambda^{m_1} \varphi_2^*(0) + \sum_{r=0}^{m_1} \lambda^{m_1-r} \left[\sum_{l=1}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right]. \end{array} \right.$$

Applying formula (1.20) we get (for $k \in \mathbb{Z}_{m_1+2}^{m_2+1}$)

$$\begin{aligned} y_1(k) &= \lambda^{k-(m_1+1)} y_1(m_1+1) \\ &\quad + \sum_{r=m_1+1}^{k-1} \lambda^{k-1-r} \left[\lambda^r \varphi_2^*(0) + b_{12}^{*1} \lambda^{r-m_1} \varphi_2^*(0) + \sum_{l=2}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right] \\ &= \lambda^{k-m_1-1} \left(\lambda^{m_1+1} \varphi_1^*(0) + (m_1+1) \lambda^{m_1} \varphi_2^*(0) \right. \\ &\quad \left. + \sum_{r=0}^{m_1} \lambda^{m_1-r} \left[\sum_{l=1}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right] \right) \\ &\quad + \sum_{r=m_1+1}^{k-1} \lambda^{k-1-r} \left[\lambda^r \varphi_2^*(0) + b_{12}^{*1} \lambda^{r-m_1} \varphi_2^*(0) + \sum_{l=2}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right] \\ &= \lambda^k \varphi_1^*(0) + (m_1+1) \lambda^{k-1} \varphi_2^*(0) + \sum_{r=0}^{m_1} \lambda^{k-1-r} \left[\sum_{l=1}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right] \\ &\quad + (k - m_1 - 1) \lambda^{k-1} \varphi_2^*(0) \\ &\quad + \sum_{r=m_1+1}^{k-1} \lambda^{k-1-r} \left[b_{12}^{*1} \lambda^{r-m_1} \varphi_2^*(0) + \sum_{l=2}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right] \\ &= \lambda^k \varphi_1^*(0) + k \lambda^{k-1} \varphi_2^*(0) + \sum_{r=0}^{k-1} \lambda^{k-1-r} \left[\sum_{l=2}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right] \\ &\quad + b_{12}^{*1} \left[\sum_{r=0}^{m_1} \lambda^{k-1-r} \varphi_2^*(r - m_1) + (k-1-m_1) \lambda^{k-1-m_1} \varphi_2^*(0) \right]. \end{aligned} \quad (2.71)$$

Now we solve equation (2.69) for $k \in \mathbb{Z}_{m_2+1}^{m_3}$ with initial data deduced from (2.71),

i.e., we consider the problem.

$$\left\{ \begin{array}{l} y_1(k+1) = \lambda y_1(k) + \lambda^k \varphi_2^*(0) + \sum_{l=1}^2 b_{12}^{*l} \lambda^{k-m_l} \varphi_2^*(0) + \sum_{l=3}^n b_{12}^{*l} \varphi_2^*(k-m_l) \\ k \in \mathbb{Z}_{m_2+1}^{m_3}, \\ y_1(m_2+1) = \lambda^{m_2+1} \varphi_1^*(0) + (m_2+1) \lambda^{m_2} \varphi_2^*(0) + \sum_{r=0}^{m_2} \lambda^{m_2-r} \left[\sum_{l=2}^n b_{12}^{*l} \varphi_2^*(r-m_l) \right] \\ + b_{12}^{*1} \left[\sum_{r=0}^{m_1} \lambda^{m_2-r} \varphi_2^*(r-m_1) + (m_2-m_1) \lambda^{m_2-m_1} \varphi_2^*(0) \right]. \end{array} \right.$$

Applying formula (1.20) yields (for $k \in \mathbb{Z}_{m_2+2}^{m_3+1}$)

$$\begin{aligned} y_1(k) &= \lambda^{k-(m_2+1)} y_1(m_2+1) \\ &+ \sum_{r=m_2+1}^{k-1} \lambda^{k-1-r} \left[\lambda^r \varphi_2^*(0) + \sum_{l=1}^2 b_{12}^{*l} \lambda^{r-m_l} \varphi_2^*(0) + \sum_{l=3}^n b_{12}^{*l} \varphi_2^*(r-m_l) \right] \\ &= \lambda^{k-m_2-1} \left(\lambda^{m_2+1} \varphi_1^*(0) + (m_2+1) \lambda^{m_2} \varphi_2^*(0) + \sum_{r=0}^{m_2} \lambda^{m_2-r} \left[\sum_{l=2}^n b_{12}^{*l} \varphi_2^*(r-m_l) \right] \right. \\ &\quad \left. + b_{12}^{*1} \left[\sum_{r=0}^{m_1} \lambda^{m_2-r} \varphi_2^*(r-m_1) + (m_2-m_1) \lambda^{m_2-m_1} \varphi_2^*(0) \right] \right) \\ &+ \sum_{r=m_2+1}^{k-1} \lambda^{k-1-r} \left[\lambda^r \varphi_2^*(0) + \sum_{l=1}^2 b_{12}^{*l} \lambda^{r-m_l} \varphi_2^*(0) + \sum_{l=3}^n b_{12}^{*l} \varphi_2^*(r-m_l) \right] \\ &= \lambda^k \varphi_1^*(0) + (m_2+1) \lambda^{k-1} \varphi_2^*(0) + \sum_{r=0}^{m_2} \lambda^{k-1-r} \left[\sum_{l=2}^n b_{12}^{*l} \varphi_2^*(r-m_l) \right] \\ &\quad + b_{12}^{*1} \left[\sum_{r=0}^{m_1} \lambda^{k-1-r} \varphi_2^*(r-m_1) + (m_2-m_1) \lambda^{k-1-m_1} \varphi_2^*(0) \right] \\ &\quad + (k-1-m_2) \lambda^{k-1} \varphi_2^*(0) \\ &\quad + \sum_{r=m_2+1}^{k-1} \lambda^{k-1-r} \left[\sum_{l=1}^2 b_{12}^{*l} \lambda^{r-m_l} \varphi_2^*(0) + \sum_{l=3}^n b_{12}^{*l} \varphi_2^*(r-m_l) \right] \\ &= \lambda^k \varphi_1^*(0) + k \lambda^{k-1} \varphi_2^*(0) + \sum_{r=0}^{k-1} \lambda^{k-1-r} \left[\sum_{l=3}^n b_{12}^{*l} \varphi_2^*(r-m_l) \right] \\ &\quad + b_{12}^{*1} \left[\sum_{r=0}^{m_1} \lambda^{k-1-r} \varphi_2^*(r-m_1) + (k-1-m_1) \lambda^{k-1-m_1} \varphi_2^*(0) \right] \\ &\quad + b_{12}^{*2} \left[\sum_{r=0}^{m_2} \lambda^{k-1-r} \varphi_2^*(r-m_2) + (k-1-m_2) \lambda^{k-1-m_2} \varphi_2^*(0) \right]. \end{aligned} \tag{2.72}$$

From (2.70), (2.71) and (2.72) we deduce that expected form of the solution of the initial problem for $k \in \mathbb{Z}_{m_s-1+1}^{m_s}$ with initial data derived from the solution of previous equation for $k \in \mathbb{Z}_{m_s-2+1}^{m_s-1}$ is

$$\begin{aligned}
 y_1(k) &= \lambda^k \varphi_1^*(0) + k \lambda^{k-1} \varphi_2^*(0) + \sum_{r=0}^{k-1} \lambda^{k-1-r} \left[\sum_{l=s}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right] \\
 &+ \sum_{l=1}^{s-1} b_{12}^{*l} \left[\sum_{r=0}^{m_l} \lambda^{k-1-r} \varphi_2^*(r - m_l) + (k-1-m_l) \lambda^{k-1-m_l} \varphi_2^*(0) \right] \\
 &\text{if } k \in \mathbb{Z}_{m_{s-1}+2}^{m_s+1}.
 \end{aligned} \tag{2.73}$$

We solve equation (2.69) for $k \in \mathbb{Z}_{m_s+1}^{m_s+1}$ with initial data deduced from (2.73), i.e., we consider the problem

$$\left\{ \begin{array}{l}
 y_1(k+1) = \lambda y_1(k) + \lambda^k \varphi_2^*(0) + \sum_{l=1}^s b_{12}^{*l} \lambda^{k-m_l} \varphi_2^*(0) + \sum_{l=s+1}^n b_{12}^{*l} \varphi_2^*(k - m_l), \\
 k \in \mathbb{Z}_{m_s+1}^{m_s+1}, \\
 y_1(m_s+1) = \lambda^{m_s+1} \varphi_1^*(0) + (m_s+1) \lambda^{m_s} \varphi_2^*(0) + \sum_{r=0}^{m_s} \lambda^{m_s-r} \left[\sum_{l=s}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right] \\
 + \sum_{l=1}^{s-1} b_{12}^{*l} \left[\sum_{r=0}^{m_l} \lambda^{m_s-r} \varphi_2^*(r - m_l) + (m_s - m_l) \lambda^{m_s-m_l} \varphi_2^*(0) \right].
 \end{array} \right.$$

Applying formula (1.20) yields (for $k \in \mathbb{Z}_{m_s+2}^{m_s+1+1}$)

$$\begin{aligned}
 y_1(k) &= \lambda^{k-(m_s+1)} y_1(m_s+1) + \sum_{r=m_s+1}^{k-1} \lambda^{k-1-r} \left[\lambda^r \varphi_2^*(0) + \sum_{l=1}^s b_{12}^{*l} \lambda^{r-m_l} \varphi_2^*(0) \right. \\
 &\quad \left. + \sum_{l=s+1}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right] \\
 &= \lambda^{k-m_s-1} \left(\lambda^{m_s+1} \varphi_1^*(0) + (m_s+1) \lambda^{m_s} \varphi_2^*(0) + \sum_{r=0}^{m_s} \lambda^{m_s-r} \left[\sum_{l=s}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right] \right. \\
 &\quad \left. + \sum_{l=1}^{s-1} b_{12}^{*l} \left[\sum_{r=0}^{m_l} \lambda^{m_s-r} \varphi_2^*(r - m_l) + (m_s - m_l) \lambda^{m_s-m_l} \varphi_2^*(0) \right] \right) \\
 &\quad + \sum_{r=m_s+1}^{k-1} \lambda^{k-1-r} \left[\lambda^r \varphi_2^*(0) + \sum_{l=1}^s b_{12}^{*l} \lambda^{r-m_l} \varphi_2^*(0) + \sum_{l=s+1}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right] \\
 &= \lambda^k \varphi_1^*(0) + (m_s+1) \lambda^{k-1} \varphi_2^*(0) + \sum_{r=0}^{m_s} \lambda^{k-1-r} \left[\sum_{l=s}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right] \\
 &\quad + \sum_{l=1}^{s-1} b_{12}^{*l} \left[\sum_{r=0}^{m_l} \lambda^{k-1-r} \varphi_2^*(r - m_l) + (m_s - m_l) \lambda^{k-1-m_l} \varphi_2^*(0) \right] \\
 &\quad + (k-1-m_s) \lambda^{k-1} \varphi_2^*(0) + \sum_{r=m_s+1}^{k-1} \lambda^{k-1-r} \left[\sum_{l=1}^s b_{12}^{*l} \lambda^{r-m_l} \varphi_2^*(0) \right. \\
 &\quad \left. + \sum_{l=s+1}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right]
 \end{aligned}$$

$$\begin{aligned}
 &= \lambda^k \varphi_1^*(0) + k \lambda^{k-1} \varphi_2^*(0) + \sum_{r=0}^{k-1} \lambda^{k-1-r} \left[\sum_{l=s+1}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right] \\
 &\quad + \sum_{r=0}^{m_s} \lambda^{k-1-r} b_{12}^{*s} \varphi_2^*(r - m_s) \\
 &\quad + b_{12}^{*1} \left[\sum_{r=0}^{m_1} \lambda^{k-1-r} \varphi_2^*(r - m_1) + (m_s - m_1) \lambda^{k-1-m_1} \varphi_2^*(0) \right] \\
 &\quad + b_{12}^{*2} \left[\sum_{r=0}^{m_2} \lambda^{k-1-r} \varphi_2^*(r - m_2) + (m_s - m_2) \lambda^{k-1-m_2} \varphi_2^*(0) \right] \\
 &\quad + \dots \\
 &\quad + b_{12}^{*s-1} \left[\sum_{r=0}^{m_{s-1}} \lambda^{k-1-r} \varphi_2^*(r - m_{s-1}) + (m_s - m_{s-1}) \lambda^{k-1-m_{s-1}} \varphi_2^*(0) \right] \\
 &\quad + (k-1-m_s) \left[\lambda^{k-1-m_1} b_{12}^{*1} \varphi_2^*(0) + \lambda^{k-1-m_2} b_{12}^{*2} \varphi_2^*(0) \right. \\
 &\quad \left. + \dots + \lambda^{k-1-m_{s-1}} b_{12}^{*s-1} \varphi_2^*(0) + \lambda^{k-1-m_s} b_{12}^{*s} \varphi_2^*(0) \right] \\
 &= \lambda^k \varphi_1^*(0) + k \lambda^{k-1} \varphi_2^*(0) + \sum_{r=0}^{k-1} \lambda^{k-1-r} \left[\sum_{l=s+1}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right] \\
 &\quad + b_{12}^{*1} \left[\sum_{r=0}^{m_1} \lambda^{k-1-r} \varphi_2^*(r - m_1) + (k-1-m_1) \lambda^{k-1-m_1} \varphi_2^*(0) \right] \\
 &\quad + b_{12}^{*2} \left[\sum_{r=0}^{m_2} \lambda^{k-1-r} \varphi_2^*(r - m_2) + (k-1-m_2) \lambda^{k-1-m_2} \varphi_2^*(0) \right] \\
 &\quad + \dots \\
 &\quad + b_{12}^{*s-1} \left[\sum_{r=0}^{m_{s-1}} \lambda^{k-1-r} \varphi_2^*(r - m_{s-1}) + (k-1-m_{s-1}) \lambda^{k-1-m_{s-1}} \varphi_2^*(0) \right] \\
 &\quad + b_{12}^{*s} \left[\sum_{r=0}^{m_s} \lambda^{k-1-r} \varphi_2^*(r - m_s) + (k-1-m_s) \lambda^{k-1-m_s} \varphi_2^*(0) \right] \\
 &= \lambda^k \varphi_1^*(0) + k \lambda^{k-1} \varphi_2^*(0) + \sum_{r=0}^{k-1} \lambda^{k-1-r} \left[\sum_{l=s+1}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right] \\
 &\quad + \sum_{l=1}^s b_{12}^{*l} \left[\sum_{r=0}^{m_l} \lambda^{k-1-r} \varphi_2^*(r - m_l) + (k-1-m_l) \lambda^{k-1-m_l} \varphi_2^*(0) \right]. \tag{2.74}
 \end{aligned}$$

In the end, we solve equation (2.69) for $k \in \mathbb{Z}_{m_n+1}^\infty$ with initial data deduced

from (2.74), i.e., we consider the problem

$$\begin{cases} y_1(k+1) = \lambda y_1(k) + \lambda^k \varphi_2^*(0) + \sum_{l=1}^n b_{12}^{*l} \lambda^{k-m_l} \varphi_2^*(0) & \text{if } k \in \mathbb{Z}_{m_n+1}^\infty, \\ y_1(m_n+1) = \lambda^{m_n+1} \varphi_1^*(0) + (m_n+1) \lambda^{m_n} \varphi_2^*(0) + \sum_{r=0}^{m_n} \lambda^{m_n-r} b_{12}^{*n} \varphi_2^*(r-m_n) \\ \quad + \sum_{l=1}^{n-1} b_{12}^{*l} \left[\sum_{r=0}^{m_l} \lambda^{m_n-r} \varphi_2^*(r-m_l) + (m_n-m_l) \lambda^{m_n-m_l} \varphi_2^*(0) \right]. \end{cases}$$

Applying formula (1.20) yields (for $k \in \mathbb{Z}_{m_n+2}^\infty$)

$$\begin{aligned} y_1(k) &= \lambda^{k-(m_n+1)} y_1(m_n+1) + \sum_{r=m_n+1}^{k-1} \lambda^{k-1-r} \left[\lambda^r \varphi_2^*(0) + \sum_{l=1}^n b_{12}^{*l} \lambda^{r-m_l} \varphi_2^*(0) \right] \\ &= \lambda^{k-m_n-1} \left(\lambda^{m_n+1} \varphi_1^*(0) + (m_n+1) \lambda^{m_n} \varphi_2^*(0) + \sum_{r=0}^{m_n} \lambda^{m_n-r} b_{12}^{*n} \varphi_2^*(r-m_n) \right. \\ &\quad \left. + \sum_{l=1}^{n-1} b_{12}^{*l} \left[\sum_{r=0}^{m_l} \lambda^{m_n-r} \varphi_2^*(r-m_l) + (m_n-m_l) \lambda^{m_n-m_l} \varphi_2^*(0) \right] \right) \\ &\quad + \sum_{r=m_n+1}^{k-1} \lambda^{k-1-r} \left[\lambda^r \varphi_2^*(0) + \sum_{l=1}^n b_{12}^{*l} \lambda^{r-m_l} \varphi_2^*(0) \right] \\ &= \lambda^k \varphi_1^*(0) + (m_n+1) \lambda^{k-1} \varphi_2^*(0) + \sum_{r=0}^{m_n} \lambda^{k-1-r} b_{12}^{*n} \varphi_2^*(r-m_n) \\ &\quad + \sum_{l=1}^{n-1} b_{12}^{*l} \left[\sum_{r=0}^{m_l} \lambda^{k-1-r} \varphi_2^*(r-m_l) + (m_n-m_l) \lambda^{k-1-m_l} \varphi_2^*(0) \right] \\ &\quad + (k-1-m_n) \lambda^{k-1} \varphi_2^*(0) + \sum_{r=m_n+1}^{k-1} \lambda^{k-1-r} \left[\sum_{l=1}^n b_{12}^{*l} \lambda^{r-m_l} \varphi_2^*(0) \right] \\ &= \lambda^k \varphi_1^*(0) + k \lambda^{k-1} \varphi_2^*(0) + \sum_{r=0}^{m_n} \lambda^{k-1-r} b_{12}^{*n} \varphi_2^*(r-m_n) \\ &\quad + b_{12}^{*1} \left[\sum_{r=0}^{m_1} \lambda^{k-1-r} \varphi_2^*(r-m_1) + (m_n-m_1) \lambda^{k-1-m_1} \varphi_2^*(0) \right] \\ &\quad + b_{12}^{*2} \left[\sum_{r=0}^{m_2} \lambda^{k-1-r} \varphi_2^*(r-m_2) + (m_n-m_2) \lambda^{k-1-m_2} \varphi_2^*(0) \right] \\ &\quad + \dots \\ &\quad + b_{12}^{*n-1} \left[\sum_{r=0}^{m_{n-1}} \lambda^{k-1-r} \varphi_2^*(r-m_{n-1}) + (m_n-m_{n-1}) \lambda^{k-1-m_{n-1}} \varphi_2^*(0) \right] \\ &\quad + (k-1-m_n) \left[\lambda^{k-1-m_1} b_{12}^{*1} \varphi_2^*(0) + \lambda^{k-1-m_2} b_{12}^{*2} \varphi_2^*(0) \right. \\ &\quad \left. + \dots + \lambda^{k-1-m_{n-1}} b_{12}^{*n-1} \varphi_2^*(0) + \lambda^{k-1-m_n} b_{12}^{*n} \varphi_2^*(0) \right] \end{aligned}$$

$$\begin{aligned}
 &= \lambda^k \varphi_1^*(0) + k \lambda^{k-1} \varphi_2^*(0) \\
 &\quad + b_{12}^{*1} \left[\sum_{r=0}^{m_1} \lambda^{k-1-r} \varphi_2^*(r - m_1) + (k - 1 - m_1) \lambda^{k-1-m_1} \varphi_2^*(0) \right] \\
 &\quad + b_{12}^{*2} \left[\sum_{r=0}^{m_2} \lambda^{k-1-r} \varphi_2^*(r - m_2) + (k - 1 - m_2) \lambda^{k-1-m_2} \varphi_2^*(0) \right] \\
 &\quad + \dots \\
 &\quad + b_{12}^{*n-1} \left[\sum_{r=0}^{m_{n-1}} \lambda^{k-1-r} \varphi_2^*(r - m_{n-1}) + (k - 1 - m_{n-1}) \lambda^{k-1-m_{n-1}} \varphi_2^*(0) \right] \\
 &\quad + b_{12}^{*n} \left[\sum_{r=0}^{m_n} \lambda^{k-1-r} \varphi_2^*(r - m_n) + (k - 1 - m_n) \lambda^{k-1-m_n} \varphi_2^*(0) \right] \\
 &= \lambda^k \varphi_1^*(0) + k \lambda^{k-1} \varphi_2^*(0) + \sum_{l=1}^n b_{12}^{*l} \left[\sum_{r=0}^{m_l} \lambda^{k-1-r} \varphi_2^*(r - m_l) \right. \\
 &\quad \left. + (k - 1 - m_l) \lambda^{k-1-m_l} \varphi_2^*(0) \right]. \tag{2.75}
 \end{aligned}$$

Summing up all particular cases (2.70)–(2.75), we get

$$y_1(k) = \left\{ \begin{array}{l} \varphi_1^*(k) \quad \text{if } k \in \mathbb{Z}_{-m_n}^0, \\ \lambda^k \varphi_1^*(0) + k\lambda^{k-1} \varphi_2^*(0) + \sum_{r=0}^{k-1} \lambda^{k-1-r} \left[\sum_{l=1}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right] \\ \quad \text{if } k \in \mathbb{Z}_1^{m_1+1}, \\ \lambda^k \varphi_1^*(0) + k\lambda^{k-1} \varphi_2^*(0) + \sum_{r=0}^{k-1} \lambda^{k-1-r} \left[\sum_{l=2}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right] \\ \quad + b_{12}^{*1} \left[\sum_{r=0}^{m_1} \lambda^{k-1-r} \varphi_2^*(r - m_1) \right. \\ \quad \left. + (k-1-m_1) \lambda^{k-1-m_1} \varphi_2^*(0) \right] \quad \text{if } k \in \mathbb{Z}_{m_1+2}^{m_2+1}, \\ \lambda^k \varphi_1^*(0) + k\lambda^{k-1} \varphi_2^*(0) + \sum_{r=0}^{k-1} \lambda^{k-1-r} \left[\sum_{l=3}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right] \\ \quad + b_{12}^{*1} \left[\sum_{r=0}^{m_1} \lambda^{k-1-r} \varphi_2^*(r - m_1) \right. \\ \quad \left. + (k-1-m_1) \lambda^{k-1-m_1} \varphi_2^*(0) \right] \\ \quad + b_{12}^{*2} \left[\sum_{r=0}^{m_2} \lambda^{k-1-r} \varphi_2^*(r - m_2) \right. \\ \quad \left. + (k-1-m_2) \lambda^{k-1-m_2} \varphi_2^*(0) \right] \quad \text{if } k \in \mathbb{Z}_{m_2+2}^{m_3+1}, \\ \dots \\ \lambda^k \varphi_1^*(0) + k\lambda^{k-1} \varphi_2^*(0) + \sum_{r=0}^{k-1} \lambda^{k-1-r} \left[\sum_{l=s+1}^n b_{12}^{*l} \varphi_2^*(r - m_l) \right] \\ \quad + \sum_{l=1}^s b_{12}^{*l} \left[\sum_{r=0}^{m_l} \lambda^{k-1-r} \varphi_2^*(r - m_l) \right. \\ \quad \left. + (k-1-m_l) \lambda^{k-1-m_l} \varphi_2^*(0) \right] \quad \text{if } k \in \mathbb{Z}_{m_s+2}^{m_{s+1}+1}, \\ \dots \\ \lambda^k \varphi_1^*(0) + k\lambda^{k-1} \varphi_2^*(0) + \sum_{l=1}^n b_{12}^{*l} \left[\sum_{r=0}^{m_l} \lambda^{k-1-r} \varphi_2^*(r - m_l) \right. \\ \quad \left. + (k-1-m_l) \lambda^{k-1-m_l} \varphi_2^*(0) \right] \quad \text{if } k \in \mathbb{Z}_{m_n+2}^\infty. \end{array} \right. \quad (2.76)$$

Formulas (2.64) and (2.65) are consequences of (2.76), (2.68).

□

3 DIMENSION OF THE SET OF SOLUTIONS

Since all the possible cases of the planar system (1.1) with weak delay have been analysed, we are ready to formulate results concerning the dimension of the space of solutions of (1.1) assuming that initial conditions (1.4) are variable. Although the case $b_{11}^{*l} = b_{22}^{*l} = b_{12}^{*l} = b_{21}^{*l} = 0$ does not lead to a weakly delayed system and is excluded by (1.3), for completeness of analysis we incorporate such possibility in our analysis as well (such a case can be considered as a degenerated weakly delayed system). Before formulation we remark that if an assumption in the following theorem is assumed to be valid for a fixed index $l \in \{1, 2, \dots, n\}$, it is easy to see that it must be valid for all indices $l = 1, 2, \dots, n$.

Theorem 3.0.6. *Let (1.1) be a weakly delayed system and (2.1) have both roots different from zero and $l \in \{1, 2, \dots, n\}$ be fixed. Then the space of solutions, being initially $2(m_n + 1)$ -dimensional, becomes on $\mathbb{Z}_{m_n+2}^\infty$ only*

1) $(m_n + 2)$ -dimensional if equation (2.1) has

- a) two real distinct roots and $(b_{12}^{*l})^2 + (b_{21}^{*l})^2 > 0$.
- b) a double real root, $b_{12}^{*l}b_{21}^{*l} = 0$ and $(b_{12}^{*l})^2 + (b_{21}^{*l})^2 > 0$.
- c) a double real root and $b_{12}^{*l}b_{21}^{*l} \neq 0$.

2) 2-dimensional if equation (2.1) has

- a) two real distinct roots and $b_{12}^{*l} = b_{21}^{*l} = 0$.
- b) a pair of complex conjugate roots.
- c) a double real root and $b_{12}^{*l} = b_{21}^{*l} = 0$.

Proof. We will carefully go through all the theorems considered (Theorem 2.1.1–Theorem 2.1.5) adding the case of a pair of complex conjugate roots and our conclusion will hold at least on $\mathbb{Z}_{m_n+2}^\infty$ (some of the statements hold on a larger interval).

a) Analysing the statement of Theorem 2.1.1 (the case (2.2) of two real distinct roots), we obtain the following subcases:

- a1) If $b_{11}^{*l} = b_{22}^{*l} = b_{21}^{*l} = 0$, $b_{12}^{*l} \neq 0$, then the dimension of the space of solutions on $\mathbb{Z}_{m_n+2}^\infty$ equals $m_n + 2$ since the last formula in (2.13) uses only $m_n + 2$ arbitrary parameters

$$\varphi_1^*(0), \varphi_2^*(-m_n), \varphi_2^*(-m_n + 1), \dots, \varphi_2^*(0).$$

- a2)** If $b_{11}^{*l} = b_{22}^{*l} = b_{12}^{*l} = 0$, $b_{21}^{*l} \neq 0$, then the dimension of the space of solutions on $\mathbb{Z}_{m_n+2}^\infty$ equals $m_n + 2$ since the last formula in (2.14) uses only $m_n + 2$ arbitrary parameters

$$\varphi_1^*(-m_n), \varphi_1^*(-m_n + 1), \dots, \varphi_1^*(0), \varphi_2^*(0).$$

- a3)** If $b_{12}^{*l} = b_{21}^{*l} = 0$, then $b_{11}^{*l} = b_{22}^{*l} = 0$ and Theorem 2.1.1 is not applicable. The dimension of the space of solutions on $\mathbb{Z}_{m_n+2}^\infty$ equals 2 since the solution is determined only by 2 arbitrary parameters

$$\varphi_1^*(0), \varphi_2^*(0).$$

This means that all the cases considered are covered by conclusions 1) a) and 2) a) of Theorem 3.0.6.

- b)** In the case (2.3) of two complex conjugate roots, we have $b_{11}^{*l} = b_{22}^{*l} = b_{12}^{*l} = b_{21}^{*l} = 0$ (i.e. we deal not with a weakly delayed system, as noted above) and the formula (2.29) uses only 2 arbitrary parameters

$$\varphi_1^*(0), \varphi_2^*(0)$$

for every $k \in \mathbb{Z}_1^\infty$. This is covered by case 2) b) of Theorem 3.0.6.

- c)** Analysing the statement of Theorem 2.1.3 and Theorem 2.1.4 (the case (2.4) of a double real root), we obtain the following subcases:

- c1)** If $b_{21}^{*l} = 0$, $b_{12}^{*l} \neq 0$, then the dimension of the space of solutions on $\mathbb{Z}_{m_n+2}^\infty$ equals $m_n + 2$ since the last formula in (2.35) uses only $m_n + 2$ arbitrary parameters

$$\varphi_1^*(0), \varphi_2^*(-m_n), \varphi_2^*(-m_n + 1), \dots, \varphi_2^*(0).$$

- c2)** If $b_{12}^{*l} = 0$, $b_{21}^{*l} \neq 0$, then the dimension of the space of solutions on $\mathbb{Z}_{m_n+2}^\infty$ equals $m_n + 2$ since the last formula in (2.36) uses only $m_n + 2$ arbitrary parameters

$$\varphi_1^*(-m_n), \varphi_1^*(-m_n + 1), \dots, \varphi_1^*(0), \varphi_2^*(0).$$

- c3)** If $b_{12}^{*l} = b_{21}^{*l} = 0$ (degenerated weakly delayed system), then the dimension of the space of solutions on $\mathbb{Z}_{m_n+2}^\infty$ equals 2 and solutions are determined only by 2 arbitrary parameters

$$\varphi_1^*(0), \varphi_2^*(0).$$

- c4)** If $b_{12}^{*l}b_{21}^{*l} \neq 0$, then the dimension of the space of solutions on $\mathbb{Z}_{m_n+2}^\infty$ equals $m_n + 2$ since the last formula in (2.42) uses only $m_n + 2$ arbitrary parameters

$$C(-m_n), C(-m_n + 1), \dots, C(0), \varphi_1^*(0)$$

where

$$C(k) := \left[\varphi_1^*(k) + \frac{b_{12}^{*l}}{b_{11}^{*l}} \varphi_2^*(k) \right], \quad k \in \mathbb{Z}_{-m_n}^0.$$

The parameter $\varphi_2^*(0)$ cannot be seen as independent since it depends on the independent parameters $\varphi_1^*(0)$ and $C(0)$.

All the cases considered are covered by conclusions 1) b), 1) c) and 2) c) of Theorem 3.0.6.

- d)** Analysing the statement of Theorem 2.1.5 (the case (2.5) of a double real root), we obtain the following subcases:

- d1)** If $b_{11}^{*l} = b_{22}^{*l} = b_{21}^{*l} = 0$, $b_{12}^{*l} \neq 0$, then the dimension of the space of solutions on $\mathbb{Z}_{m_n+2}^\infty$ equals $m_n + 2$ since the last formula in (2.64) uses only $m_n + 2$ arbitrary parameters

$$\varphi_1^*(0), \varphi_2^*(-m_n), \varphi_2^*(-m_n + 1), \dots, \varphi_2^*(0)$$

and the last formula in (2.65) provides no new information.

- d2)** If $b_{11}^{*l} = b_{22}^{*l} = b_{21}^{*l} = b_{12}^{*l} = 0$ (degenerated weakly delayed system), then the dimension of the space of solutions on $\mathbb{Z}_{m_n+2}^\infty$ equals 2 since solutions are determined only by 2 arbitrary parameters

$$\varphi_1^*(0), \varphi_2^*(0).$$

Both cases are covered by conclusions 1) b) and 2) c) of Theorem 3.0.6.

Since there are no cases other than the above cases a)–d), the proof is finished. \square

Theorem 3.0.6 can be formulated simply as

Theorem 3.0.7. *Let (1.1) be a weakly delayed system and let (2.1) have both roots different from zero. Then the space of solutions, being initially $2(m_n + 1)$ -dimensional, is on $\mathbb{Z}_{m_n+2}^\infty$ only*

- 1) $(m_n + 2)$ -dimensional if $(b_{12}^{*l})^2 + (b_{21}^{*l})^2 > 0$.
- 2) 2-dimensional if $b_{12}^{*l} = b_{21}^{*l} = 0$.

We omit the proofs of the following two theorems since, again, are much the same as those of Theorems 2.1.1–2.1.5.

Theorem 3.0.8. *Let (1.1) be a weakly delayed system and let (2.1) have a simple root $\lambda = 0$. Then the space of solutions, being initially $2(m_n + 1)$ -dimensional, is either $(m_n + 1)$ -dimensional or 1-dimensional on $\mathbb{Z}_{m_n+2}^\infty$.*

Theorem 3.0.9. *Let (1.1) be a weakly delayed system and let (2.1) have a double root $\lambda = 0$. Then the space of solutions, being initially $2(m_n + 1)$ -dimensional, turns into a 0-dimensional space on $\mathbb{Z}_{m_n+2}^\infty$, namely, into the zero solution.*

4 DISCUSSION OF STABILITY

In this chapter we use the explicit formulas derived in Chapter 3 for stability analysis of linear system (1.1).

Define a norm of a 2×2 matrix $A = \{a_{ij}\}_{i,j=1}^2$ as

$$\|A\| = \max\{|a_{11}| + |a_{12}|, |a_{21}| + |a_{22}|\}$$

and, for 2×1 vectors $x = (x_1, x_2)^T$, an induced vector norm

$$\|x\| = \max\{|x_1|, |x_2|\}.$$

For a discrete vector $\psi: \mathbb{Z}_{-m_n}^0 \rightarrow \mathbb{R}^2$ we define

$$\|\psi\|_{m_n} := \max\{\|\psi(-m_n)\|, \|\psi(-m_n + 1)\|, \dots, \|\psi(0)\|\}.$$

Now we define the stability of the zero solution of linear system (1.1)

$$x(k+1) = Ax(k) + \sum_{l=1}^n B^l x_l(k - m_l)$$

where $k \in \mathbb{Z}_{-m_n}^\infty$, B^l are 2×2 constant matrices.

Definition 4.0.10. *The zero solution $x(k) = 0$, $k \in \mathbb{Z}_{-m_n}^\infty$ of (1.1) is said to be*

- a) *Stable if, given $\varepsilon > 0$ and $k_0 \geq 0$, there exists $\delta = \delta(\varepsilon, k_0)$ such that $\varphi(k)$, $k \in \mathbb{Z}_{k_0 - m_n}^{k_0}$, $\|\varphi\|_{m_n} < \delta$ implies $\|x(k, k_0, \varphi)\| < \varepsilon$ for all $k \geq k_0$, uniformly stable if δ may be chosen independently of k_0 , unstable if it is not stable;*
- b) *Asymptotically stable if it is stable and $\lim_{k \rightarrow \infty} \|x(k)\| = 0$;*
- c) *Conditionally stable (conditionally asymptotically stable) if it is stable (asymptotically stable) under the condition that a subspace P of the space all initial data with $\dim P$ satisfying*

$$1 < \dim P < 2(m_n + 1)$$

is fixed.

It is easy to see that

$$\max\{\|\varphi^*(0)\|, \|\Phi_2(0)\|, \|\Phi_2(-1)\|, \dots, \|\Phi_2(-m_l)\|\} \leq \|\varphi^*\|_{m_n}.$$

Assume

$$q = \max\{|\lambda_1|, |\lambda_2|\} < 1.$$

Lemma 4.0.11. *For every $k \in \mathbb{Z}_{m_n+2}^\infty$, the following inequality*

$$(k - 1 - m_l)q^{k-1-m_l} \leq M := -\frac{1}{q^{\frac{1}{\ln q}} \ln q} \quad (4.1)$$

holds.

Proof. Define an auxiliary function

$$f(x) = (x - 1 - m_l)q^{x-1-m_l} \quad (4.2)$$

and determine its maximum. The first derivate of the function (4.2) is

$$f'(x) = q^{x-1-m_l} + (x - 1 - m_l)q^{x-1-m_l} \ln q.$$

From the equation

$$q^{x-1-m_l} + (x - 1 - m_l)q^{x-1-m_l} \ln q = 0$$

we obtain a stationary point

$$x_0 = 1 + m_l - \frac{1}{\ln q}.$$

The second derivative of the function $f(x)$ is

$$\begin{aligned} f''(x) &= q^{x-1-m_l} \ln q + q^{x-1-m_l} \ln q + (x - 1 - m_l)q^{x-1-m_l} \ln^2 q \\ &= q^{x-1-m_l} \ln q \left[2 + (x - 1 - m_l) \ln q \right]. \end{aligned}$$

At the stationary point x_0 , we get

$$f''\left(1 + m_l - \frac{1}{\ln q}\right) = q^{-\frac{1}{\ln q}} \ln q. \quad (4.3)$$

The value of (4.3) is negative for $q < 1$ and, at the stationary point, the function $f(x)$ assumes its maximum. Since $f(x_0) = M$, formula (4.1) holds. \square

In the following Theorems 4.0.12–4.0.27, we prove the stability, asymptotic stability, conditional asymptotic stability, and conditional stability of system (2.6), (2.7). Tracing the statements of these theorems we conclude that it is sufficient to perform stability investigation only on the interval $k \in \mathbb{Z}_{m_n+2}^\infty$. Therefore we omit the technical details connected with investigating the stability on $k \in \mathbb{Z}_0^{m_n+1}$.

Theorem 4.0.12. *If Theorem 2.1.1 holds and $q < 1$, then the zero solution of (1.1) is stable.*

Proof. Let the case I) be valid. We show that the zero solution is stable (we assume $k \in \mathbb{Z}_{m_n+2}^\infty$).

$$\begin{aligned}
 \|y(k)\| &\leq \|\Lambda_1^k\| \|\varphi^*(0)\| \\
 &+ \left\| \sum_{l=1}^n b_{12}^{*l} \left[\sum_{r=0}^{m_l} \lambda_1^{k-1-r} \Phi_2(r - m_l) + \Phi_2(0) \sum_{r=m_l+1}^{k-1} \lambda_1^{k-1-r} \lambda_2^{r-m_l} \right] \right\| \\
 &\leq \|\Lambda_1^k\| \|\varphi^*(0)\| \\
 &+ \sum_{l=1}^n |b_{12}^{*l}| \left[\sum_{r=0}^{m_l} |\lambda_1|^{k-1-r} \|\Phi_2(r - m_l)\| + \|\Phi_2(0)\| \sum_{r=m_l+1}^{k-1} |\lambda_1|^{k-1-r} |\lambda_2|^{r-m_l} \right] \\
 &\leq \max\{|\lambda_1|^k; |\lambda_2|^k\} \|\varphi^*(0)\| \\
 &+ \sum_{l=1}^n |b_{12}^{*l}| \left[\sum_{r=0}^{m_l} |\lambda_1|^{k-1-r} \|\Phi_2(r - m_l)\| + \|\Phi_2(0)\| \sum_{r=m_l+1}^{k-1} |\lambda_1|^{k-1-r} |\lambda_2|^{r-m_l} \right] \\
 &\leq q^k \|\varphi^*\|_{m_n} + \sum_{l=1}^n |b_{12}^{*l}| \left[\sum_{r=0}^{m_l} q^{k-1-r} \|\varphi^*\|_{m_n} + \|\varphi^*\|_{m_n} \sum_{r=m_l+1}^{k-1} q^{k-1-m_l} \right] \\
 &\leq \left[q^k + \sum_{l=1}^n |b_{12}^{*l}| \left(q^{k-1-m_l} \frac{1 - q^{m_l+1}}{1 - q} + (k - 1 - m_l) q^{k-1-m_l} \right) \right] \|\varphi^*\|_{m_n} \\
 &\leq \left[1 + \sum_{l=1}^n |b_{12}^{*l}| \left(\frac{1 - q^{m_l+1}}{1 - q} + M \right) \right] \|\varphi^*\|_{m_n}. \tag{4.4}
 \end{aligned}$$

We set

$$\begin{aligned}
 E &:= \left[1 + \sum_{l=1}^n |b_{12}^{*l}| \left(\frac{1 - q^{m_l}}{1 - q} + M \right) \right], \\
 \delta &:= \frac{\varepsilon}{E}.
 \end{aligned}$$

This equality implies

$$\|y(k)\| \leq E \|\varphi^*\|_{m_n} < \varepsilon, k \in \mathbb{Z}_{m_n+2}^\infty$$

if

$$\|\varphi^*\|_{m_n} < \delta.$$

If the case II) holds, the proof can be performed similarly. \square

Moreover, we can improve Theorem 4.0.12 since the inequality $q < 1$ guarantees even asymptotic stability.

Theorem 4.0.13. *If Theorem 2.1.1 holds and $q < 1$, then the zero solution of (1.1) is asymptotically stable.*

Proof. Let the case I) be valid. By Theorem 4.0.12, the zero solution of (1.1) is stable. We show that

$$\lim_{k \rightarrow \infty} \|y(k)\| = 0$$

for every initial problem (1.1) is asymptotically stable for $k \in \mathbb{Z}_{m_n+2}^\infty$. From the inequalities obtained in the proof of Theorem 4.0.12, we get

$$\begin{aligned} & \lim_{k \rightarrow \infty} \|y(k)\| \\ & \leq \lim_{k \rightarrow \infty} \left[q^k + \sum_{l=1}^n |b_{12}^{*l}| \left(q^{k-1-m_l} \frac{1-q^{m_l+1}}{1-q} + (k-1-m_l)q^{k-1-m_l} \right) \right] \|\varphi^*\|_{m_n} \\ & \leq \lim_{k \rightarrow \infty} q^k \|\varphi^*\|_{m_n} + \lim_{k \rightarrow \infty} \left(\sum_{l=1}^n |b_{12}^{*l}| q^{k-1-m_l} \frac{1-q^{m_l+1}}{1-q} \right) \|\varphi^*\|_{m_n} \\ & \quad + \lim_{k \rightarrow \infty} \left(\sum_{l=1}^n |b_{12}^{*l}| (k-1-m_l) q^{k-1-m_l} \right) \|\varphi^*\|_{m_n} = 0 \end{aligned}$$

where the last limit was computed by L' Hospital rule. If the case II) holds, the proof can be performed similarly. \square

Theorem 4.0.14. *If Theorem 2.1.3 holds and $q < 1$, then the zero solution of (1.1) is asymptotically stable.*

Proof. We show that the zero solution is stable for the case $b_{21}^{*l} = 0$, for $k \in \mathbb{Z}_{m_n+2}^\infty$. We will proceed in a way similar to that of the proof of Theorem 4.0.12. Then,

$$\begin{aligned} \|y(k)\| & \leq \|\Lambda_3^k\| \|\varphi^*(0)\| \\ & \quad + \left\| \sum_{l=1}^n b_{12}^{*l} \left[\sum_{r=0}^{m_l} \lambda^{k-1-r} \Phi_2(r-m_l) + (k-1-m_l) \lambda^{k-1-m_l} \Phi_2(0) \right] \right\| \\ & \leq \|\Lambda_3^k\| \|\varphi^*(0)\| \\ & \quad + \sum_{l=1}^n |b_{12}^{*l}| \left[\sum_{r=0}^{m_l} |\lambda|^{k-1-r} \|\Phi_2(r-m_l)\| + (k-1-m_l) |\lambda|^{k-1-m_l} \|\Phi_2(0)\| \right] \\ & \leq |\lambda|^k \|\varphi^*(0)\| \\ & \quad + \sum_{l=1}^n |b_{12}^{*l}| \left[\sum_{r=0}^{m_l} |\lambda|^{k-1-r} \|\Phi_2(r-m_l)\| + (k-1-m_l) |\lambda|^{k-1-m_l} \|\Phi_2(0)\| \right] \\ & \leq q^k \|\varphi^*\|_{m_n} + \sum_{l=1}^n |b_{12}^{*l}| \left[\sum_{r=0}^{m_l} q^{k-1-r} \|\varphi^*\|_{m_n} + (k-1-m_l) q^{k-1-m_l} \|\varphi^*\|_{m_n} \right] \\ & \leq \left[q^k + \sum_{l=1}^n |b_{12}^{*l}| \left(q^{k-1-m_l} \frac{1-q^{m_l+1}}{1-q} + (k-1-m_l) q^{k-1-m_l} \right) \right] \|\varphi^*\|_{m_n} \end{aligned}$$

$$\leq \left[1 + \sum_{l=1}^n |b_{12}^{*l}| \left(\frac{1 - q^{m_l+1}}{1 - q} + M \right) \right] \|\varphi^*\|_{m_n}.$$

We obtain the same inequality (4.4) as in the proof of Theorem 4.0.12. The rest of the proof of stability copies that of Theorem 4.0.12. Asymptotic stability can be proved by a scheme similar to that of the proof of Theorem 4.0.13. If $b_{21}^{*l} = 0$, the proof can be performed similarly. \square

Theorem 4.0.15. *If Theorem 2.1.4 holds and $q < 1$, then the zero solution of (1.1) is asymptotically stable.*

Proof. For $\|y(k)\|$, $k \in \mathbb{Z}_{m_n+2}^\infty$ we get

$$\begin{aligned} \|y(k)\| &\leq \|\Lambda_3^k\| \|\varphi^*(0)\| \\ &+ \left\| \sum_{l=1}^n \left[\sum_{r=0}^{m_l} \lambda^{k-1-r} \Phi_l^*(r - m_l) + (k - 1 - m_l) \lambda^{k-1-m_l} \Phi_l^*(0) \right] \right\| \\ &\leq \|\Lambda_3^k\| \|\varphi^*(0)\| \\ &+ \sum_{l=1}^n \left[\sum_{r=0}^{m_l} |\lambda|^{k-1-r} \|\Phi_l^*(r - m_l)\| + (k - 1 - m_l) |\lambda|^{k-1-m_l} \|\Phi_l^*(0)\| \right] \\ &\leq |\lambda|^k \|\varphi^*(0)\| \\ &+ \sum_{l=1}^n \left[\sum_{r=0}^{m_l} |\lambda|^{k-1-r} \|\Phi_l^*(r - m_l)\| + (k - 1 - m_l) |\lambda|^{k-1-m_l} \|\Phi_l^*(0)\| \right] \\ &\leq q^k \|\varphi^*\|_{m_n} + \sum_{l=1}^n \left[\sum_{r=0}^{m_l} q^{k-1-r} \|\Phi_l^*\|_{m_n} + (k - 1 - m_l) q^{k-1-m_l} \|\Phi_l^*\|_{m_n} \right] \\ &\leq q^k \|\varphi^*\|_{m_n} + \left[\sum_{l=1}^n \left(q^{k-1-m_l} \frac{1 - q^{m_l+1}}{1 - q} + (k - 1 - m_l) q^{k-1-m_l} \right) \right] \|\Phi^*\|_{m_n} \\ &\leq \|\varphi^*\|_{m_n} + \left[\sum_{l=1}^n \left(\frac{1 - q^{m_l+1}}{1 - q} + M \right) \right] \|\varphi^*\|_{m_n} \\ &\leq \left[1 + \sum_{l=1}^n \left(\frac{1 - q^{m_l+1}}{1 - q} + M \right) \right] \|\varphi^*\|_{m_n}. \end{aligned} \tag{4.5}$$

From (4.5), it is easy to see that the zero solution of (1.1) is stable. The proof of asymptotic stability can be made similarly to that of Theorem 4.0.13. \square

Lemma 4.0.16. *Let $q \in (0, 1)$, then, for every $k \in \mathbb{Z}_{m_n+2}^\infty$, the following inequality*

$$q^k + kq^{k-1} \leq N := q^{-\frac{q \ln q + 1}{\ln q} - 1} \left(-\frac{1}{\ln q} \right) \tag{4.6}$$

holds.

Proof. Define an auxiliary function

$$f(x) = q^x + xq^{x-1} \quad (4.7)$$

and determine its maximum. The first derivate of (4.7) is

$$f'(x) = q^x \ln q + q^{x-1} + xq^{x-1} \ln q.$$

From the equation

$$q^x \ln q + q^{x-1} + xq^{x-1} \ln q = 0,$$

we obtain a stationary point

$$x_0 = -\frac{q \ln q + 1}{\ln q}.$$

The second derivative of the function $f(x)$ is

$$\begin{aligned} f''(x) &= q^x \ln^2 q + q^{x-1} \ln q + q^{x-1} \ln q + xq^{x-1} \ln^2 q \\ &= q^{x-1} \ln q \left(q \ln q + 2 + x \ln q \right). \end{aligned}$$

At the stationary point x_0 , we have

$$f''\left(-\frac{q \ln q + 1}{\ln q}\right) = q^{-\frac{q \ln q + 1}{\ln q} - 1} \ln q. \quad (4.8)$$

The value of (4.8) is negative for $q < 1$ and, at the stationary point, the function $f(x)$ assumes its maximum. Since $f(x_0) = N$, formula (4.6) holds. \square

Theorem 4.0.17. *If Theorem 2.1.5 holds and $q < 1$, then the zero solution of (1.1) is asymptotically stable.*

Proof. First, we show that the zero solution of (1.1) is stable. For $\|y_1(k)\|$, $k \in \mathbb{Z}_{m_n+2}^\infty$ we get

$$\begin{aligned} \|y_1(k)\| &\leq |\lambda|^k \|\varphi_1^*(0)\| + k|\lambda|^{k-1} \|\varphi_2^*(0)\| \\ &\quad + \left\| \sum_{l=1}^n b_{12}^{*l} \left[\sum_{r=0}^{m_l} \lambda^{k-1-r} \varphi_2^*(r - m_l) + (k-1-m_l) \lambda^{k-1-m_l} \varphi_2^*(0) \right] \right\| \\ &\leq q^k \|\varphi^*\|_{m_n} + kq^{k-1} \|\varphi^*\|_{m_n} \\ &\quad + \sum_{l=1}^n |b_{12}^{*l}| \left[\sum_{r=0}^{m_l} |\lambda|^{k-1-r} \|\varphi^*\|_{m_n} + (k-1-m_l) |\lambda|^{k-1-m_l} \|\varphi^*\|_{m_n} \right] \end{aligned}$$

$$\begin{aligned}
 &\leq \left[q^k + kq^{k-1} \right. \\
 &\quad \left. + \sum_{l=1}^n |b_{12}^{*l}| \left(q^{k-1-m_l} \frac{1-q^{m_l+1}}{1-q} + (k-1-m_l)q^{k-1-m_l} \right) \right] \|\varphi^*\|_{m_n} \\
 &\leq \left[N + \sum_{l=1}^n |b_{12}^{*l}| \left(\frac{1-q^{m_l+1}}{1-q} + M \right) \right] \|\varphi^*\|_{m_n} \tag{4.9}
 \end{aligned}$$

and, for $\|y_2(k)\|$, $k \in \mathbb{Z}_{m_n+2}^\infty$, we get

$$\|y_2(k)\| \leq |\lambda|^k \|\varphi_2^*(0)\| \leq q^k \|\varphi^*\|_{m_n} \leq \|\varphi^*\|_{m_n}, k \in \mathbb{Z}_{m_n+2}^\infty. \tag{4.10}$$

From (4.9), (4.10) it is easy to see that the zero solution of (1.1) is stable. The proof of asymptotic stability can be made similarly to that of Theorem 4.0.13. \square

Theorem 4.0.18. *If Theorem 2.1.1 holds and $|\lambda_1| = 1, |\lambda_2| \leq q < 1$, then the zero solution of (1.1) is stable.*

Proof. We show that the zero solution of (1.1) is stable for $k \in \mathbb{Z}_{m_n+2}^\infty$. For the case I), we get

$$\begin{aligned}
 \|y(k)\| &\leq \|\Lambda_1^k\| \|\varphi^*(0)\| \\
 &\quad + \left\| \sum_{l=1}^n b_{12}^{*l} \left[\sum_{r=0}^{m_l} \lambda_1^{k-1-r} \Phi_2(r-m_l) + \Phi_2(0) \sum_{r=m_l+1}^{k-1} \lambda_1^{k-1-r} \lambda_2^{r-m_l} \right] \right\| \\
 &\leq \|\Lambda_1^k\| \|\varphi^*(0)\| \\
 &\quad + \sum_{l=1}^n |b_{12}^{*l}| \left[\sum_{r=0}^{m_l} |\lambda_1|^{k-1-r} \|\Phi_2(r-m_l)\| + \|\Phi_2(0)\| \sum_{r=m_l+1}^{k-1} |\lambda_1|^{k-1-r} |\lambda_2|^{r-m_l} \right] \\
 &\leq \max\{|\lambda_1|^k; |\lambda_2|^k\} \|\varphi^*(0)\| \\
 &\quad + \sum_{l=1}^n |b_{12}^{*l}| \left[\sum_{r=0}^{m_l} |\lambda_1|^{k-1-r} \|\Phi_2(r-m_l)\| + \|\Phi_2(0)\| \sum_{r=m_l+1}^{k-1} |\lambda_1|^{k-1-r} |\lambda_2|^{r-m_l} \right] \\
 &\leq \|\varphi^*\|_{m_n} + \sum_{l=1}^n |b_{12}^{*l}| \left[\sum_{r=0}^{m_l} \|\varphi^*\|_{m_n} + \|\varphi^*\|_{m_n} \sum_{r=m_l+1}^{k-1} q^{r-m_l} \right] \\
 &\leq \left[1 + \sum_{l=1}^n |b_{12}^{*l}| \left((m_l+1) + q \frac{1-q^{k-1-m_l}}{1-q} \right) \right] \|\varphi^*\|_{m_n} \\
 &\leq \left[1 + \sum_{l=1}^n |b_{12}^{*l}| \left((m_l+1) + \frac{q}{1-q} \right) \right] \|\varphi^*\|_{m_n}.
 \end{aligned}$$

We set

$$M_1 := \left[1 + \sum_{l=1}^n |b_{12}^{*l}| \left((m_l + 1) + \frac{q}{1-q} \right) \right],$$

$$\delta := \frac{\varepsilon}{M_1}.$$

This equality implies

$$\|y(k)\| \leq M_1 \|\varphi^*\|_{m_n} < \varepsilon, k \in \mathbb{Z}_{m_n+2}^\infty$$

if

$$\|\varphi^*\|_{m_n} < \delta.$$

In the case II), we get

$$\begin{aligned} \|y(k)\| &\leq \|\Lambda_1^k\| \|\varphi^*(0)\| \\ &+ \left\| \sum_{l=1}^n |b_{21}^{*l}| \left[\sum_{r=0}^{m_l} \lambda_2^{k-1-r} \Phi_1(r - m_l) + \Phi_1(0) \sum_{r=m_l+1}^{k-1} \lambda_1^{r-m_l} \lambda_2^{k-1-r} \right] \right\| \\ &\leq \|\Lambda_1^k\| \|\varphi^*(0)\| \\ &+ \sum_{l=1}^n |b_{21}^{*l}| \left[\sum_{r=0}^{m_l} |\lambda_2|^{k-1-r} \|\Phi_1(r - m_l)\| + \|\Phi_1(0)\| \sum_{r=m_l+1}^{k-1} |\lambda_1|^{r-m_l} |\lambda_2|^{k-1-r} \right] \\ &\leq \max\{|\lambda_1|^k; |\lambda_2|^k\} \|\varphi^*(0)\| \\ &+ \sum_{l=1}^n |b_{21}^{*l}| \left[\sum_{r=0}^{m_l} |\lambda_2|^{k-1-r} \|\Phi_1(r - m_l)\| + \|\Phi_1(0)\| \sum_{r=m_l+1}^{k-1} |\lambda_1|^{r-m_l} |\lambda_2|^{k-1-r} \right] \\ &\leq \|\varphi^*\|_{m_n} + \sum_{l=1}^n |b_{21}^{*l}| \left[\sum_{r=0}^{m_l} q^{k-1-r} \|\varphi^*\|_{m_n} + \|\varphi^*\|_{m_n} \sum_{r=m_l+1}^{k-1} q^{k-1-r} \right] \\ &\leq \left[1 + \sum_{l=1}^n |b_{21}^{*l}| \sum_{r=0}^{k-1} q^{k-1-r} \right] \|\varphi^*\|_{m_n} \\ &\leq \left[1 + \sum_{l=1}^n |b_{21}^{*l}| \frac{1-q^k}{1-q} \right] \|\varphi^*\|_{m_n} \\ &\leq \left[1 + \sum_{l=1}^n |b_{21}^{*l}| \frac{1}{1-q} \right] \|\varphi^*\|_{m_n}. \end{aligned}$$

We set

$$M_2 := \left[1 + \sum_{l=1}^n |b_{21}^{*l}| \frac{1}{1-q} \right],$$

$$\delta := \frac{\varepsilon}{M_2}.$$

This equality implies

$$\|y(k)\| \leq M_2 \|\varphi^*\|_{m_n} < \varepsilon, k \in \mathbb{Z}_{m_n+2}^\infty$$

if

$$\|\varphi^*\|_{m_n} < \delta.$$

□

Theorem 4.0.19. *If Theorem 2.1.1 holds and $|\lambda_1| \leq q < 1, |\lambda_2| = 1$, then the zero solution of (1.1) is stable.*

Proof. In this case, we carry out the proof similarly to that of Theorem 4.0.18

□

Theorem 4.0.20. *If Theorem 2.1.1 holds, the case I) occurs, $|\lambda_1| \leq q < 1, |\lambda_2| \geq 1$ and $\varphi_2^*(0) = 0$, then the zero solution of (1.1) is conditionally asymptotically stable.*

Proof. In this case, $\varphi^*(0) = (\varphi_1^*(0), 0)^T$ and $\Phi_2(0) = (\varphi_2^*(0), 0)^T = (0, 0)$, we perform the proof similarly to that of Theorem 4.0.18 and Theorem 4.0.13. For $k \in \mathbb{Z}_{m_n+2}^\infty$, we get

$$\begin{aligned} \|y(k)\| &\leq \|\Lambda_1^k \varphi^*(0)\| \\ &+ \left\| \sum_{l=1}^n b_{12}^{*l} \left[\sum_{r=0}^{m_l} \lambda_1^{k-1-r} \Phi_2(r - m_l) + \Phi_2(0) \sum_{r=m_l+1}^{k-1} \lambda_1^{k-1-r} \lambda_2^{r-m_l} \right] \right\| \\ &\leq \left\| \begin{pmatrix} \lambda_1^k & 0 \\ 0 & \lambda_2^k \end{pmatrix} \begin{pmatrix} \varphi_1^*(0) \\ 0 \end{pmatrix} \right\| \\ &+ \sum_{l=1}^n |b_{12}^{*l}| \left[\sum_{r=0}^{m_l} |\lambda_1|^{k-1-r} \|\Phi_2(r - m_l)\| + \|\Phi_2(0)\| \sum_{r=m_l+1}^{k-1} |\lambda_1|^{k-1-r} |\lambda_2|^{r-m_l} \right] \\ &\leq |\lambda_1|^k \|\varphi_1^*(0)\| \\ &+ \sum_{l=1}^n |b_{12}^{*l}| \left[\sum_{r=0}^{m_l} |\lambda_1|^{k-1-r} \|\Phi_2(r - m_l)\| \right] \\ &\leq q^k \|\varphi^*\|_{m_n} + \sum_{l=1}^n |b_{12}^{*l}| \left[\sum_{r=0}^{m_l} q^{k-1-r} \|\varphi^*\|_{m_n} \right] \\ &\leq \left[q^k + \sum_{l=1}^n |b_{12}^{*l}| \sum_{r=0}^{m_l} q^{k-1-r} \right] \|\varphi^*\|_{m_n} \\ &\leq \left[q^k + \sum_{l=1}^n |b_{12}^{*l}| \left(q^{k-1-m_l} \frac{1 - q^{m_l+1}}{1 - q} \right) \right] \|\varphi^*\|_{m_n} \end{aligned}$$

$$= q^k \left[1 + \sum_{l=1}^n |b_{12}^{*l}| \frac{q^{-m_l-1} - 1}{1 - q} \right] \|\varphi^*\|_{m_n}.$$

Now, it is easy to see that

$$\lim_{k \rightarrow \infty} \|y(k)\| = 0.$$

□

Theorem 4.0.21. *If Theorem 2.1.1 holds, the case II) occurs, $|\lambda_2| \leq q < 1$, $|\lambda_1| \geq 1$ and $\varphi_1^*(0) = 0$, then the zero solution of (1.1) is conditionally asymptotically stable.*

Proof. In this case, $\varphi^*(0) = (0, \varphi_2^*(0))^T$ and $\Phi_1(0) = (0, \varphi_1^*(0))^T = (0, 0)$, we perform the proof similarly to that of Theorem 4.0.20. For $k \in \mathbb{Z}_{m_n+2}^\infty$, we get

$$\begin{aligned} \|y(k)\| &\leq \|\Lambda_1^k \varphi^*(0)\| \\ &+ \left\| \sum_{l=1}^n b_{21}^{*l} \left[\sum_{r=0}^{m_l} \lambda_2^{k-1-r} \Phi_1(r - m_l) + \Phi_1(0) \sum_{r=m_l+1}^{k-1} \lambda_1^{r-m_l} \lambda_2^{k-1-r} \right] \right\| \\ &\leq \left\| \begin{pmatrix} \lambda_1^k & 0 \\ 0 & \lambda_2^k \end{pmatrix} \begin{pmatrix} 0 \\ \varphi_2^*(0) \end{pmatrix} \right\| \\ &+ \sum_{l=1}^n |b_{21}^{*l}| \left[\sum_{r=0}^{m_l} |\lambda_2|^{k-1-r} \|\Phi_1(r - m_l)\| + \|\Phi_1(0)\| \sum_{r=m_l+1}^{k-1} |\lambda_1|^{r-m_l} |\lambda_2|^{k-1-r} \right] \\ &\leq |\lambda_2|^k \|\varphi_2^*(0)\| \\ &+ \sum_{l=1}^n |b_{21}^{*l}| \left[\sum_{r=0}^{m_l} |\lambda_2|^{k-1-r} \|\Phi_1(r - m_l)\| \right] \\ &\leq q^k \|\varphi^*\|_{m_n} + \sum_{l=1}^n |b_{21}^{*l}| \left[\sum_{r=0}^{m_l} q^{k-1-r} \|\varphi^*\|_{m_n} \right]. \end{aligned}$$

We obtain the same inequality as in the proof of Theorem 4.0.20. The rest of the proof copies that of Theorem 4.0.20. □

Theorem 4.0.22. *If Theorem 2.1.1 holds, the case I) occurs, $|\lambda_1| = 1$, $|\lambda_2| > 1$ and $\varphi_2^*(0) = 0$, then the zero solution of (1.1) is conditionally stable.*

Proof. In this case, $\varphi^*(0) = (\varphi_1^*(0), 0)^T$ and $\Phi_2(0) = (\varphi_2^*(0), 0)^T = (0, 0)$ we, perform the proof similarly to that of Theorem 4.0.20. For $k \in \mathbb{Z}_{m_n+2}^\infty$, we get

$$\begin{aligned} \|y(k)\| &\leq \|\Lambda_1^k \varphi^*(0)\| \\ &+ \left\| \sum_{l=1}^n b_{12}^{*l} \left[\sum_{r=0}^{m_l} \lambda_1^{k-1-r} \Phi_2(r - m_l) + \Phi_2(0) \sum_{r=m_l+1}^{k-1} \lambda_1^{k-1-r} \lambda_2^{r-m_l} \right] \right\| \end{aligned}$$

$$\begin{aligned}
 &\leq \left\| \begin{pmatrix} \lambda_1^k & 0 \\ 0 & \lambda_2^k \end{pmatrix} \begin{pmatrix} \varphi_1^*(0) \\ 0 \end{pmatrix} \right\| \\
 &\quad + \sum_{l=1}^n |b_{12}^{*l}| \left[\sum_{r=0}^{m_l} |\lambda_1|^{k-1-r} \|\Phi_2(r - m_l)\| + \|\Phi_2(0)\| \sum_{r=m_l+1}^{k-1} |\lambda_1|^{k-1-r} |\lambda_2|^{r-m_l} \right] \\
 &\leq |\lambda_1|^k \|\varphi_1^*(0)\| \\
 &\quad + \sum_{l=1}^n |b_{12}^{*l}| \left[\sum_{r=0}^{m_l} |\lambda_1|^{k-1-r} \|\Phi_2(r - m_l)\| \right] \\
 &\leq \|\varphi^*\|_{m_n} + \sum_{l=1}^n |b_{12}^{*l}| \left[\sum_{r=0}^{m_l} \|\varphi^*\|_{m_n} \right] \\
 &\leq \left[1 + \sum_{l=1}^n |b_{12}^{*l}| (m_l + 1) \right] \|\varphi^*\|_{m_n}.
 \end{aligned}$$

We set

$$\begin{aligned}
 M_3 &:= 1 + \sum_{l=1}^n |b_{12}^{*l}| (m_l + 1), \\
 \delta &:= \frac{\varepsilon}{M_3}.
 \end{aligned}$$

This equality implies

$$\|y(k)\| \leq M_3 \|\varphi^*\|_{m_n} < \varepsilon, k \in \mathbb{Z}_{m_n+2}^\infty$$

if

$$\|\varphi^*\|_{m_n} < \delta.$$

□

Theorem 4.0.23. *If Theorem 2.1.1 holds, the case II) occurs, $|\lambda_2| = 1, |\lambda_1| > 1$ and $\varphi_1^*(0) = 0$, then the zero solution of (1.1) is conditionally stable.*

Proof. We perform the proof similarly to that of Theorem 4.0.22. □

Theorem 4.0.24. *If Theorem 2.1.3 holds, the case $b_{21}^{*l} = 0$ occurs, $|\lambda| = 1$ and $\varphi_2^*(0) = 0$, then the zero solution of (1.1) is conditionally stable.*

Proof. In this case, $\varphi^*(0) = (\varphi_1^*(0), 0)^T$ and $\Phi_2(0) = (0, 0)^T$, we perform the proof similarly to that of Theorem 4.0.14 and, for $k \in \mathbb{Z}_{m_n+2}^\infty$, we get

$$\begin{aligned}
 \|y(k)\| &\leq \|\Lambda_3^k \varphi^*(0)\| \\
 &\quad + \left\| \sum_{l=1}^n b_{12}^{*l} \left[\sum_{r=0}^{m_l} \lambda^{k-1-r} \Phi_2(r - m_l) + (k - 1 - m_l) \lambda^{k-1-m_l} \Phi_2(0) \right] \right\|
 \end{aligned}$$

$$\begin{aligned}
 &\leq \left\| \begin{pmatrix} \lambda^k & 0 \\ 0 & \lambda^k \end{pmatrix} \begin{pmatrix} \varphi_1^*(0) \\ 0 \end{pmatrix} \right\| \\
 &\quad + \sum_{l=1}^n |b_{12}^{*l}| \left[\sum_{r=0}^{m_l} |\lambda|^{k-1-r} \|\Phi_2(r - m_l)\| + (k-1-m_l) |\lambda|^{k-1-m_l} \|\Phi_2(0)\| \right] \\
 &\leq \|\varphi_1^*(0)\| + \sum_{l=1}^n |b_{12}^{*l}| \left[\sum_{r=0}^{m_l} \|\varphi^*\|_{m_n} + (k-1-m_l) \|\Phi_2(0)\| \right] \\
 &\leq \left[1 + \sum_{l=1}^n |b_{12}^{*l}| (m_l + 1) \right] \|\varphi^*\|_{m_n}.
 \end{aligned}$$

We set

$$\begin{aligned}
 M_4 &:= 1 + \sum_{l=1}^n |b_{12}^{*l}| (m_l + 1), \\
 \delta &:= \frac{\varepsilon}{M_4}.
 \end{aligned}$$

This equality implies

$$\|y(k)\| \leq M_4 \|\varphi^*\|_{m_n} < \varepsilon, k \in \mathbb{Z}_{m_n+2}^\infty$$

if

$$\|\varphi^*\|_{m_n} < \delta.$$

□

Theorem 4.0.25. *If Theorem 2.1.3 holds, the case $b_{12}^{*l} = 0$ occurs, $|\lambda| = 1$ and $\varphi_1^*(0) = 0$, then the zero solution of (1.1) is conditionally stable.*

Proof. We perform the proof similarly to that of Theorem 4.0.24. □

Theorem 4.0.26. *If Theorem 2.1.4 holds and $|\lambda| = 1$ and $\varphi_1^*(0) = \varphi_2^*(0) = 0$, then the zero solution of (1.1) is conditionally stable.*

Proof. Conditional stability can be proved using a scheme similar to that of the proof of Theorem 4.0.24. □

Theorem 4.0.27. *If Theorem 2.1.5 holds and $|\lambda| = 1$ and $\varphi_2^*(0) = 0$, then the zero solution of (1.1) is conditionally stable.*

Proof. We show that the zero solution of (1.1) is conditionally stable. For $\|y_1(k)\|$, $k \in \mathbb{Z}_{m_n+2}^\infty$, we get

$$\begin{aligned}
 \|y_1(k)\| &\leq |\lambda|^k \|\varphi_1^*(0)\| + k |\lambda|^{k-1} \|\varphi_2^*(0)\| \\
 &\quad + \left\| \sum_{l=1}^n b_{12}^{*l} \left[\sum_{r=0}^{m_l} \lambda^{k-1-r} \varphi_2^*(r - m_l) + (k-1-m_l) \lambda^{k-1-m_l} \varphi_2^*(0) \right] \right\|
 \end{aligned}$$

$$\begin{aligned} &\leq \|\varphi_1^*(0)\| + \sum_{l=1}^n |b_{12}^{*l}| \left[\sum_{r=0}^{m_l} \|\varphi^*\|_{m_n} \right] \\ &\leq 1 + \sum_{l=1}^n |b_{12}^{*l}| (m_l + 1) \|\varphi^*\|_{m_n}, \end{aligned} \tag{4.11}$$

and, for $\|y_2(k)\|$, $k \in \mathbb{Z}_{m_n+2}^\infty$, we get

$$\|y_2(k)\| \leq |\lambda|^k \|\varphi_2^*(0)\| = 0, k \in \mathbb{Z}_{m_n+2}^\infty. \tag{4.12}$$

From (4.11), (4.12) it is easy to see that the zero solution of (1.1) is conditionally stable. \square

5 CONCLUSIONS

To our best knowledge, weakly delayed systems were first defined in [24] for systems of linear delayed differential systems with constant coefficients and, in [15], for planar linear discrete systems with a single delay (in these papers such systems are called systems with a weak delay). The weakly delayed systems analyzed in this paper can be simplified and their solutions can be found in explicit analytical forms (results obtained and published in [10] generalize those in [15] and [9]). Consequently, analytical forms of solutions can be used directly to solve several problems for weakly delayed systems, e.g., problems of asymptotical behavior of their solutions, stability problems, boundary-value problems, and some problems of control theory.

REFERENCES

- [1] Agarwal R. P.; Bohner M., Grace S. R.; O'Regan D. *Discrete oscillation theory*. Hindawi Publishing Corporation, New York, 2005. xiv+961 pp. ISBN: 977-5945-19-4.
- [2] Appleby J.; Guzowska M.; Rodkina A. *Preservation of positivity in the solution of discretized stochastic differential equation*. Mathematical Science, preprint archive, Dublin City University, School of Mathematics, MS-07-21, (2007), 1–15.
- [3] Baštinec J.; Diblík J. *One case of appearance of positive solutions of delayed discrete equations*,. Applications of Mathematics, 48 (2003), 429–436. ISSN: 0862-7940.
- [4] Baštinec J.; Diblík J. *Subdominant positive solutions of the discrete equation $\Delta u(k+n) = -p(k)u(k)$* . Abstr. Appl. Anal., 2004:6 (2004), 461–470. ISSN: 1085–3375, e-ISSN: 1687–0409.
- [5] Baštinec J.; Diblík J.; Šmarda Z. *Existence of positive solutions of discrete linear equations with a single delay*. Difference Equ. Appl., 16, Issue 9 (2010), 1047–1056. ISSN: 1563-5120 (electronic) 1023-6198 (paper).
- [6] Bereketoğlu H.; Huseynov A. *Convergence of solutions of nonhomogeneous linear difference systems with delays*. Acta Appl. Math., 110 (2010), No 1, 259–269.
- [7] Čermák J. *On the related asymptotics of delay differential and difference equations*. Dynam. Systems Appl., 14, No 3–4, (2005), 419–429. ISSN: 1056–2176.
- [8] Čermák J. *Asymptotic bounds for linear difference systems*. Adv. Difference Equ., Volume 2010 (2010), Article ID 182696, 14 pages, doi:10.1155/2010/182696.
- [9] Diblík, J.; Halfarová, H. *Explicit general solution of planar linear discrete systems with constant coefficient and weak delays*. Adv. Difference Equ. 2013, Art. number: 50, doi:10.1186/1687-1847-2013-50, 1–29.
- [10] Diblík, J.; Halfarová, H. *General explicit solution of planar weakly delayed linear discrete systems and pasting its solutions*. Abstr. Appl. Anal. 2014, doi:10.1155/2014/627295, 1–37.
- [11] Diblík J.; Hlavičková I. *Combination of Liapunov and retract methods in the investigation of the asymptotic behavior of solutions of systems of discrete equations*. Dynam. Systems Appl., 18, Issue 3–4 (2009), 507–537.

- [12] Diblík J.; Hlavičková I. *Asymptotic behavior of solutions of delayed difference equations*. Abstr. Appl. Anal., vol. 2011, Article ID 671967, 24 pages, 2011. doi:10.1155/2011/671967.
- [13] Diblík, J.; Khusainov, D. Ya. *Representation of solutions of linear discrete systems with constant coefficients and pure delay*. Adv. Difference Equ. 2006, Art. ID 80825, DOI 10.1155/ADE/2006/80825, 1–13. (doi:10.1155/ADE)
- [14] Diblík J.; Khusainov D. Ya. *Representation of solutions of discrete delayed system $x(k+1) = Ax(k) + Bx(k-m) + f(k)$ with commutative matrices*. J. Math. Anal. Appl. 318 (2006), no. 1, 63–76.
- [15] Diblík, J.; Khusainov, D. Ya.; Šmarda, Z. *Construction of the general solution of planar linear discrete systems with constant coefficients and weak delay*. Adv. Difference Equ. 2009, Art. ID 784935, 18 pages, doi:10.1155/2009/784935.
- [16] Diblík J.; Khusainov D. Ya.; Grytsay I.V.; Šmarda Z. *Stability of nonlinear autonomous quadratic discrete systems in the critical case*. Discrete Dynam. Nat. Soc., vol. 2010, Article ID 539087, 23 pages, 2010. doi:10.1155/2010/539087.
- [17] Diblík J.; Růžičková M.; Schmeidel E., Zbąszyniak M. *Weighted asymptotically periodic solutions of linear Volterra difference equations*. Abstr. Appl. Anal., vol. 2011, Article ID 370982, 14 pages, 2011. doi:10.1155/2011/370982.
- [18] Diblík J.; Růžičková M.; Šmarda Z.; Šutá Z. *Asymptotic convergence of the solutions of a dynamic equation on discrete time scales*. Abstr. Appl. Anal., vol. 2011, Article ID 580750, 20 pages, doi:10.1155/2012/580750.
- [19] Elaydi, S. N. *An Introduction to Difference Equations*. Third Edition, Springer, 2005. xxii+539 pp. ISBN:0-387-23059-9.
- [20] Györi I.; Horváth L.; Reynolds D. *On asymptotically periodic solutions of linear discrete Volterra equations*, Fasc. Math., 44 (2010), 53–67.
- [21] Györi I.; Ladas G. *Oscillation theory of delay differential equations*. With applications. Oxford Mathematical Monographs. Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1991. xii+368 pp. ISBN: 0-19-853582-1.
- [22] Ivanov S.; Kipnis M.; Malygina V. *The stability cone for a difference matrix equation with two delays*. ISRN Applied Mathematics IN THE PRINT. ISSN: 2090-5564 (Print), ISSN: 2090-5572 (Online), doi:10.5402/APPMATH.

- [23] Karajani P.; Stavroulakis I.P. *Oscillation criteria for second-order delay and functional equations*. Studies of the University of Žilina, 18, No 1, (2004), 17–26. (ISSN 1336-149X)
- [24] Khusainov, D. Ya.; Benditkis, D. B.; Diblík, J. *Weak delay in systems with an aftereffect*. Funct. Differ. Equ. 9 (2002), 385-404.
- [25] Khusainov D. Ya.; Ivanov A. F.; Shuklin G. V. *On a representation of solutions of linear systems with delay*. (Russian) Differ. Uravn. 41 (2005), no. 7, 1001–1004, 1008; translation in Differ. Equ. 41 (2005), no. 7, 1054–1058
- [26] Kipnis, M.; Komissarova, D. *Stability of a delay difference system*. Adv. Difference Equ. 2006, Art. No 31409, 2006, 1–8.
- [27] Kocić V. L.; Ladas G. *Global behavior of nonlinear difference equations of higher order with applications*. Mathematics and its Applications, 256. Kluwer Academic Publishers Group, Dordrecht, 1993. xii+228 pp. ISBN: 0-7923-2286-X
- [28] Medina R. *Absolute stability of discrete-time systems with delay*. Adv. Difference Equ. 2008, Article ID 396504, 14 pages, 2008. doi:10.1155/2008/396504. (ISSN: 1687-1839, e-ISSN: 1687-1847, doi:10.1155/ADE).
- [29] Medina R.; Pituk M. *Nonoscillatory solutions of a second order difference equation of Poincaré type*. Appl. Math. Lett., 22, (2009), 679–683.
- [30] Niri K.; Stavroulakis I.P. *Oscillations of delay and difference equations*. Tatra Mt. Math. Publ., 43, (2009), 173–187.
- [31] Stavroulakis I.P. *Oscillation criteria for delay, difference and functional equations*. Funct. Differ. Equ., 11 (2004), 163–183.
- [32] Stević, S. *More on a rational recurrence relation*. Appl. Math. E-Notes 4 (2004), 80–85.
- [33] Stević, S. *On a system of difference equations*. Appl. Math. Comput. 218 (2011), 3372–3378.
- [34] Stević, S. *On the difference equation $x_n = x_{n-2}/(b_n + c_n x_{n-1} x_{n-2})$* . Appl. Math. Comput. 218 (2011), 4507–4513.
- [35] Stević, S. *On a third-order system of difference equations*. Appl. Math. Comput. 218 (2012), 7649–7654.
- [36] Stević, S. *On some solvable systems of difference equations*. Appl. Math. Comput. 218 (2012), 5010–5018.

- [37] Stević, S. *On the difference equation $x_n = x_{n-k}/(b+cx_{n-1} \dots x_{n-k})$* . *Appl. Math. Comput.* 218(2012), 6291–6296.
- [38] Stević, S.; Diblík, J.; Iričanin, B.; Šmarda, Z. *On a third-order system of difference equations with variable coefficients*. *Abstr. Appl. Anal.* vol. 2012, Article ID 508523, (2012), 1–22.
- [39] Stević, S.; Iričanin B. *Unbounded Solutions of the Difference Equation $x(n) = x(n-l)x(n-k-1)$* . *Abstr. Appl. Anal.*, Article Number: 561682, DOI: 10.1155/2011/561682, Published: 2011.